

Khalleedd ✓



Lecture:

Physiology of Brain Circulation & CSF Formation

Dr. Eyad M. Hussein

Ph.D. of Neurology

Consultant & Head of Neurology Department,
Nasser Hospital

في بداية المحاضرة حاولوا ان عندكم معلومة وبتكم تركوا فيها عتقوا الاضاءه حوالكم باختصار بدوهم نفوت بتفاصيل
لاني مشه طابق حاله صراحة ، بيقالك انو الاضاءه بتعمل constriction of the pupil وهذا بعمل
parasympathatic effect
كل هاده كلام شخصيات قوم اطفي الصواوا وادرسه كالك بتخضر فيلم ع تنقليلكس

Cerebral Blood Flow

Brain consumption about 20-25% of CO → Fifth of the cardiac output.

- **Cerebral blood flow (CBF):** is the blood supply to the brain in a given time.
- In an adult, CBF is typically **750 ml/min**.
- Regional blood supply
50-54 ml of blood per 100 g of brain tissue per min.
Normally.
- **Ischemia** results if blood flow to the brain below **18-25 ml** per 100 g per min, and **tissue death** occurs if flow decreased below **8-10 ml** per 100 g per minute.

Cerebral Blood Flow

- The adult brain weight is about 1300 – 1500 g (2% of Total Body Weight). *And take 1/5 of CO → So its a very important organ.*

- **Consumes:**

- ✓ “15-20% O₂” of the total cardiac output.

- ✓ 25% glucose

- Able to respond to variations in demand.

الدم الواصلة اليه
يعني باختصار يعرف يتأقلم على ظروف أقل من التوصل، بس لدرجة معينة طبعا

By a way called autoregulation

Brain Autoregulation


- The brain is very sensitive to over and under perfusion.
- **Cerebral autoregulation** refers to the physiological mechanisms that maintain blood flow at an appropriate level during changes in blood pressure.

مجموعة الأشياء التي يتحكم بها autoregulation


□ Physiological Mechanisms of Autoregulation:

- **Myogenic/Arterial blood pressure.** By VC+VD
- **Chemical:**
 - ✓ Arterial partial pressure of carbon dioxide **“PaCO₂”**.
 - ✓ Arterial partial pressure of oxygen **“PaO₂”**.
Decreased O₂ cuz VD
Increased O₂ cuz VC
- **Metabolic regulation.** Glucose
Hyperglycemia cuz VC
Hypoglycemia cuz VD
- **Neurogenic regulation.** Sympathetic + parasympathetic

معلومة حلوة نعرفها انهم يستخدمو الفكرة هاي في انه لو اجت على ال ICU حالة brain edema بنخدمهم على mechanical ventilation with hyperventilation علشان اعمل VC وبهيك بمنة ال brain edema



Factors Affecting Cerebral Blood Flow



Myogenic / Arterial Blood Pressure

- Arterioles ^{VD} **dilate** or ^{VC} **constrict** in response to changes in BP and ^{Intracranial pressure} ICP in order to maintain a constant ^{Cerebral (cranial) blood flow} CBF.
- The response to **lower pressure**, is arteriolar **dilation** in the brain.
- While when **blood pressure rises** they **constrict**.

بس دي عشان يكون كفو تمام تفهموا الفرق بينه ضغط الشريان وال blood flow وانو الي بيعملي انو احافظ على ال CBF وال
perfusion for tissue
في مثال انو الضغط نازل يعني الدم الي بيوصل النسيج قليل ف انا دي ازوده ف دي اوسع الشرايين
والعكس في حالة الضغط العالي

Arterial Blood Pressure

- **Cerebral Perfusion Pressure (CPP)**

- $CPP = MAP - ICP$

MAP = Mean Arterial Pressure

ICP = Intra Cranial Pressure

- **ICP** – can be measured directly.

- **MAP:** pushes blood into the brain.

- Intracranial Pressure (**ICP**).

- $MAP = \text{Diastolic pressure} + \frac{1}{3} \text{Pulse pressure}^{\text{sys-dias}}$

- **Pulse pressure** = Systolic BP – Diastolic BP.

- Eg. BP = 140/80, **MAP** = $80 + \frac{1}{3} * 60 = 100$ mmHg

Cerebral
Blood
Flow

Cerebral
Hypoxia
←
watershed syndrome
Hemodynamic changes = Dec cerebral
flow = low cerebral perfusion

Autoregulatory

Range


Headaches
BBB disruption
Edema
→

0

100

200

Mean Arterial Pressure (mmHg)




Cerebral Blood Flow

- **CBF** = Cerebral perfusion pressure(**CPP** (divided by the cerebrovascular resistance(**CVR**)).
- **CBF = CPP / CVR**
- **CPP = MAP – ICP**
- **CPP** = is normally between 70 - 90 mmHg in an adult human.
- **Increase in ICP** → decreases CBF & cerebral perfusion.

Chemical Autoregulation

- $\uparrow$ Elevation of PaCO_2 , leads to vasodilatation “High ICP”.
- $\downarrow$ Depression of PaCO_2 , leads to vasoconstriction.
- $\uparrow$ Elevation of PaO_2 , leads to vasoconstriction Treat the brain edema
“Hyperventilation used for management of brain oedema”.
- $\downarrow$ Depression of PaO_2 , “Hypoxia”: leads to vasodilatation
“High ICP”.



Arterial Blood Pressure Autoregulation

- **Baseline CBF:** 50 54 ml/100g/min
- **Critical ischemic threshold:** 20 –25 ml/100g/min
Less than it is very dangerous
- **Severe Hypertension:** High BP
 - Damage to blood brain barrier.
 - Risk of cerebral oedema and acute hypertensive encephalomalacia due to vasodilation.


المفروض مع زيادة الضغط يحصل VC

Neurogenic Autoregulation


- **Neurogenic Autoregulation:**
 - Sympathetic fibres: vasoconstriction
 - Parasympathetic fibres: vasodilatation

The Blood–Brain Barrier


- It is the barrier between cerebral capillary blood, the CSF and brain tissue.
- CSF fills the ventricles and the subarachnoid space.
- It consists of the endothelial cells of the cerebral capillaries and the choroid plexus epithelium.



Blood-Brain Barrier




Brain or spinal cord tissue



Found on choroidal plexus produce CSF

Blood-Brain Barrier

Can Pass	Cannot Pass
<ul style="list-style-type: none">Glucose (via transport proteins).Lipid-soluble substances.O₂, CO₂, alcohol, anesthetics.	<ul style="list-style-type: none">Large molecules ex. Proteins.Most antibiotic drugs.
<u>Trauma, inflammation and certain toxins can break down the BBB</u>	



Functions of the Blood–Brain Barrier

- 1. It maintains a constant environment for ^{cell bodies}neurons** in the CNS and protects the brain from ^{Metabolic and waste products.}endogenous or ^{That came from outside}exogenous toxins.
- 2. It prevents the escape of neurotransmitters** from their functional sites in the CNS into the general circulation.
- 3. Drugs penetrate the blood.**

عشان تدخل دواء الـ drugs الي مسامح الحاجات الـ penetration وتغذي الـ BBB وتكون الـ CNS

Development of BBB

- ^{الي ينولدوا على عمر سبعة او ثمانية شهور} Premature infants with ^{Bilirubin is a toxic material} hyperbilirubinemia, free bilirubin pass BBB, and may stain basal nuclei causing damage (**Kernicterus**).
^{هنا ليس حليتنا عندها nuclei وما حليتنا عندها ganglia}
^{أقلكم الفرق، الاشئ ما بسميه داخل ال CNS يطلق عليه اسم nuclei، اما الاشئ بال PNS يطلق عليه اسمك تاتي عادي}

And cause cerebral palsy with hyper kinetic type
يكون رقبته مرتفعة لفوق الدكتور مثلها تمثيل

Ganglia is nerve cells located outside the CNS

هيك خلصت جزئية الفسيو وضايك الاناتومي الله يعينكم، وأوى الله يسعدها بعثت انو بكرة
ما عنادوا ف انا بدي اسحب عليها من هاهو للمسا

Blood Supply of the Brain

Brain blood supply formed of two main systems (arteries):

1. Internal carotid system: Right & Left internal


Arise from common carotid a.

Rt from brachiocephalic a.

Lt from arch of aorta directly.

Supplies about 80% of cerebral blood flow and called ant. Circulation.

carotid arteries (ICA).



2. Vertebro-basilar system: Right & Left

Also called cerebral a.

Arise from Rt and Lt vertebral a.

Supplies about 20-25 % of the cerebral blood flow and called post. Circulation.


vertebral arteries (VA).


Rt PCA + Lt PCA ينقسمان لـ Basilar artery


Cuz it supplies cerebellum occipital lobe brain stem

PCA → posterior cerebral artery

connection between the 2 systems
internal carotid and the vertebro-basilar systems







The diagram shows a common carotid artery (CCA) bifurcating into an internal carotid artery (ICA) and an external carotid artery (ECA). A dashed line indicates the level of the C3-C4 disc. The ICA is shown continuing upwards, while the ECA branches off to the side.

I. The Internal Carotid Artery (ICA)

❖ **Origin:** arises in the neck of terminal branches of the common carotid artery CCA **(at Bifurcation)**, at the level of the upper border of the thyroid cartilage **(opposite the disc between C3 & C4)**.


Give internal and external for the face. To external and internal

❖ **Function:** the ICA supplies structures in the orbit and a large part of the brain (70- 80% of the brain blood supply).

internal carotid system


- Each ICA enters the skull through **the carotid foramen.**


- It gives **No branches** at the neck.
- branches for neck and face*




The blue matter is internal carotid artery sympathetic plexus.

Branches of the Internal Carotid Artery



- Ophthalmic artery.** 1st branch its terminal branch called central retinal artery
 - Posterior communicating artery.** PCA
Anastomosis between the vertebro-basilar system and internal carotid system.
 - Anterior choroidal artery.**
 - Anterior cerebral artery (ACA).** Medial branch
 - Middle cerebral artery (MCA).** Largest branch
Lateral branch
- 




Two terminal branches

1. Ophthalmic Artery

- **Ophthalmic artery** which enters the orbit through the optic canal with the optic nerve.
- **Supplies** the eye, orbit contents & gives **central retinal artery**.
its terminal branch
Supply the Retina (الشبكية) of the eye
- **Also supplies** the frontal sinuses, part of frontal area of scalp and dorsal part of nose.
- Its occlusion results in **Ipsilateral Blindness**.
Occlusion of central retinal artery
If the occlusion happens to the patient and not treated by anti thrombotic drugs he will face lifelong blind, cuz it has no collateral supply

Arteries and Veins of Orbit and Eyelids

Superior View




2. Posterior Communicating Artery

- The posterior communicating artery joints the **ICA** with the corresponding **posterior cerebral artery** to establish **the circle of Willis.**


➤ **It supplies:**

- Optic tract and chiasma.
- Part of diencephalon.
medially the thalamus and related structures
- Walls of third ventricle.
- A common site of **Aneurysms.** 20%
common sites for aneurysms
بأجزاء معينة من الأوعية الدموية
وكم نسبة كل وحدة منهم



Arteries of Brain

Inferior View



3. Anterior Choroidal Artery

From its name it supplies the choroidal plexus

➤ **Origin:** from the ICA just before origin of the anterior & middle cerebral arteries.

➤ **It supplies:**

- Choroid plexus of the lateral and third ventricles.
- Optic tract and optic chiasma.
- Hippocampus, tail of caudate nucleus, amygdaloid nucleus, globus pallidus and crus cerebri. .
- Lateral geniculate body.
- Posterior limb of internal capsule. Has 3 parts anterior limb, posterior limb, and genu.

Arteries of Brain [Continued]

Inferior View

Anterior communicating artery

Recurrent artery (of Heubner)

Anterior cerebral artery

Posterior communicating artery

Middle cerebral artery

Anterior choroidal artery

Optic tract

Posterior cerebral artery

Lateral geniculate body

Posterior lateral choroidal artery

Cerebral peduncle


Choroid plexus of lateral ventricle

Posterior medial choroidal artery

Medial geniculate body

Lateral ventricle

Pulvinar of thalamus



4. Anterior Cerebral Artery (ACA)

Medial branch

- It is the smaller terminal branch of the ICA.
- It runs forward and joins the ACA of the opposite side by the **anterior communicating artery** (**the commonest site of Aneurysms**).^{40%}

➤ Branches of the ACA:

1. **Cortical** branches.
2. **Central** branches (**Striate** arteries).


Territory of Anterior Cerebral Artery (ACA)

- Medial surface of the cerebral hemisphere from frontal pole to the parieto-occipital sulcus.
- Medial 1/3 of the orbital surface of the frontal lobe.
- Superior lateral surface of cerebral hemisphere (**except occipital lobe**).Supplied by PCA
- Anterior part of lentiform nucleus.
- Head of caudate nucleus.
- Supplies part of the corpus striatum and the anterior limb of the internal capsule via the **medial striate artery (of Heubner)**.

N.B.: The ACA supplies motor and sensory areas of the leg and foot “**paracentral lobule**”.Lower limb
Supero lateral lobes of the cerebrum


Arteries of Brain

Inferior View



Arteries of Brain

Medial View




5. Middle Cerebral Artery (MCA)

Lateral branch

- ❖ The lateral branch as the MCA and ACA **largest of the 2 terminal branches** of ICA.
- ❖ It pass laterally in the stem of The fissure or sulci that separate the temporal lobe from frontal and parietal lobes **the lateral sulcus.**
- ❖ It turns backwards and upwards in the posterior ramus of the lateral sulcus.
- ❖ Finally it divides into:
 - **Cortical branches.**
 - **Central branches.** Striate


Arteries of Brain

Inferior View



Arteries of Brain

Lateral View



Territory of Middle Cerebral Artery (MCA)

1. Cortical branches: supply:

- Lateral 2/3 of the orbital part of frontal lobe.
- Temporal pole and insula.
- Most of the lateral surface of cerebral hemisphere

Especially lower and middle while upper by ACA and occipital by PCA

Except:

- The upper part (superior inch): supplied by the ACA.
- Lower part of temporal lobe: supplied by the PCA.
- Occipital lobe: supplied by the PCA.
- Supplies the trunk, arm, and face areas of the cortical motor and sensory areas. *Face and upper limbs*

*Occlusion of MCA will cuz contralateral
faciobrachial hemiplegia (hemi-hypoesthesia)*

2. Central Branches: are many branches called **striate arteries**: Divided to medial and lateral according to corpus striatum

Short a. **Medial striate arteries: supply:**

- Posterior part of lentiform nucleus.
- Body of caudate nucleus.
- The most of the internal capsule (posterior limb, Except ant. Part of ant. Limb by ACA genu, and posterior part of anterior limb).


Long b. **Lateral striate arteries: supply:**

- Corpus striatum (lentiform & caudate).
- Most of the internal capsule.
- One of these arteries is large and is liable to hemorrhage it is called **the artery of cerebral hemorrhage (Charcot's artery)**.

ما نادر نيو كلينيكال حناخه ال hemorrhagic stroke و حناخه انوال intracerebral hematoma
بالية منهم ال basal nuclei وبنه هه ال ال اسم ال artery اسم ال charcots or gemorrhagic artery و هه
smooth layer very thin ال

Arteries of Brain

Frontal Section



II. The Vertebrobasilar System

A. Vertebral Artery



- **Origin:** each vertebral artery arises from the **first part of subclavian artery**. The anterior scalene muscle separates the subclavian to 3 parts.

Medial to the muscle

- Supplies structures of the brain (**20- 30% of the brain blood supply**).


- **Course:**

From **subclavian artery** → then passes upwards through the **transverse foramina** of the **upper 6 cervical vertebrae** → **foramen magnum** → they pass upward and medially in front of the **medulla oblongata** → at the lower border of pons **two vertebral arteries unite** to form the single **Basilar Artery**. Then bifurcate and give 2 post. Cerebral artery




Subclavian Artery

Right Anterior Dissection




Transverse foramen of upper 6 cervical vertebra



Branches of the Vertebral Artery

1. **The meningeal branches:** are small and supply the bone and dura in the posterior cranial fossa.
2. **PSA Posterior spinal arteries:** may arise from **the posterior inferior cerebellar artery**. *Anatomical variant*
3. **ASA Anterior spinal artery:** it unites with the other one to form the **one anterior spinal artery**.
4. **Medullary branches:** supply the ventral part of **medulla oblongata**.
5. **PICA Posterior inferior cerebellar artery.**
Largest branch

سؤال منطقي خستاله لانا وجوابه انوال vertebral artery
أقربها ال basilar artery على ال spinal cord



Posterior Inferior Cerebellar Artery (PICA)

❑ The largest branch of vertebral artery.

• It supplies the:

• Posterior part of inferior surface of cerebellum.
From its name

• Inferior cerebellar peduncle.
Connect the cerebellum with medulla oblongata

• Posteriolateral part of medulla oblongata.

• Choroid plexus of 4th ventricle.


❖ **Occlusion** of this artery leads to **lateral medullary syndrome (Wallenberg's syndrome)**.


Case: 52 y/o female with HTN and DM came to the clinic complain from sudden onset of ataxia with MRI scan we find multiple lesions and patches in cerebellum and medulla oblongata and they diagnose it mistakenly with multiple sclerosis because it was multiple lesions

النتيجة أو حدوث damage in cerebellum يقلل من قدرة المريض على التوازن والسيطرة على حركاته الدقيقة، Hypothalamus وبنابغ على ال spinal cord ال يكون حاداً ال sympathetic fibres، يعاني من انضيق التراكب ال هو ال sensation of face ال ينشأ من damage of desc. spinal nucleus اما التأثير على الجسم يحصل على السائر التالي

Arteries of Brain


Inferior View






B. Basilar Artery

- **Origin:** at the lower border of pons by the union of the two vertebral arteries.
- It ascends along the basilar groove in front of the pons.
- **Termination:** at the upper border of the pons by dividing into two terminal branches (posterior cerebral arteries).^{PCA}
- It lies medial to the abducent and oculomotor nerves.






Branches of the Basilar artery

1. **AICA** **Anterior inferior cerebellar artery:** ^{From its name} supplies the anterior part of inferior surface of the cerebellum.
2. **Internal auditory (labyrinthine) artery:** it arises either from basilar or anterior inferior cerebellar arteries. Supplies the inner ear.
Internal carotid give a branch for the eye
Basilar give for the ear
3. **Pontine (paramedian) branches:** ^{مخروطية} are small branches to supply the pons.
4. **Superior cerebellar artery:** the oculomotor nerve separates it from the posterior cerebral artery. Supplies the superior surface of cerebellum, superior cerebellar peduncle, pons and midbrain.
5. **PCA** **Posterior cerebral artery:** terminal branches.

Arteries of Brain

Inferior View




Posterior Cerebral Artery (PCA)

- It is the largest and terminal branches of the basilar artery at the upper border of the pons.
- It is separated from the superior cerebellar artery by the oculomotor nerve. The nerves lies laterally
- Connects with the ipsilateral ICA by the posterior communicating artery.

Arteries of Brain

Inferior View



Branches of Posterior Cerebral Arteries

1. Cortical Branches:

- Inferiolateral part of temporal lobe.
- Occipital lobe “Visual cortex”.
Primary and secondary

2. Central Branches:

Diencephalon and subcortical structure


- Supply parts of the thalamus, medial geniculate body, splenium of the corpus callosum, lentiform nucleus, midbrain, and pineal body.

3. Posterior Choroidal Arteries:

- supply the choroid plexus of the lateral and third ventricles.
For ICA


Arteries of Brain

Medial View



Arteries of Brain [Continued]

Inferior View




Clinical anatomy:


- The **artery of Percheron** is a **rare anatomical variation** in the brain vascularization in which a **single arterial trunk arises from the PCA** to supply **both** sides of the thalamus and midbrain.

يعني بالحالة العادية لو صار جلطة بالجزء الأيمن بيتأثر الجزء الأيمن مع ال *thalamus* حينئذ الجزء الأيمن، والعكس صحيح، بس في حالة انو انا عندي شريان واحد يغذي هاي الستمشيز، وصار فيه مشكلة في حينئذ الجانبيين مع ال *thalamus*


Normal anatomy



Anatomy with artery of Percheron



Lateral view
Coronal section through left cerebral hemisphere



Arteries


ACA

MCA

PCA


Ant choroidal

Coronal view




Medial view


Coronal section through right cerebral hemisphere




Horizontal view



Lateral view of left hemisphere



Medial view of left hemisphere




MCA

ACA

PCA

Horizontal view



Inferior view



Circle Of Willis

- ❖ **Definition:** it is an arterial anastomoses between **the vertebral and internal carotid arteries of both sides**).
- ❖ It lies in the **interpeduncular fossa** at the **base** of the **brain**
- ❖ **Function:**
 - Direct blood to other branches in cases of occlusion.
 - Connect ^{80%} carotid system with ^{20%} vertebrobasilar system.


Circle Of Willis

1. Anterior communicating artery. Between ACA
 2. Right & left of ACA.
 3. Right & left ICA.
 4. Right & left posterior communicating arteries.
 5. Right & left PCA. Post. Cerebral artery
-
6. Basilar artery. Upper part of it

9 arteries

Circle of Willis - Vessels Dissected Out

Inferior View



The Important Anastomoses Between the Cerebral Arteries

- Between anterior middle and posterior branches*
1. **Anastomoses between the branches of the cerebral arteries** on the surface of the cerebral hemisphere.
 2. **The circle of Willis:** between ICA and vertebral-basilar system.
 3. **Anastomoses between the branches of the internal and external carotid arteries:**
 - a. Anastomoses between the branches of the ICA ophthalmic artery with the facial and maxillary ECA arteries.
 - b. Anastomoses between the meningeal branches of the internal carotid artery with middle meningeal branch of maxillary artery.
ECA

Venous Drainage

A. **Superficial cerebral veins:** laceration of these vessels results in subdural hemorrhage (hematoma).

B. **Deep cerebral veins.**

C. **Dural Cerebral Sinuses.**

ال sinus هاهي بتتشأ منه انو في بعض المناطق بيصير في فراغ بيده
الطبقتيه تعود ال dura matter


Dura matter in the brain its a part of connective tissue composed 2 layers outer endostial layer and inner menengial lager
طبعا وعشاه اللام يلوهم ربط بتنا عرف انو خارج البياض زي السبايند كود يلوهم ولاير

- The veins of the brain have no muscular tissue in their thin layer.
- No valves.
- Drain into the cranial or dural venous sinuses.

Take the blood from cerebral veins and CSF

Cerebral Veins

1. **Superior cerebral veins:** drain into **the superior sagittal sinus.**
2. **Inferior cerebral veins:** drain into **the petrosal and transverse sinuses.**
3. **Middle cerebral veins:**
 - a. **Superficial middle cerebral veins:** drain into **the cavernous sinus.**
 - b. **Deep middle cerebral veins:** drain into **basal vein** → Great cerebral vein → **the straight sinus.**



3 veins

4. Basal vein (vein of Rosenthal): by union of three veins:


- **Deep middle cerebral vein.**
- **Anterior cerebral vein.**
- **Striate veins.**

- The two basal veins drain into **the great cerebral vein.**



2 veins

5. The Internal Cerebral Vein: there are two internal cerebral veins and they are formed by the union of

- **Thalamostriate vein**
- **Choroid vein.**




- The two internal cerebral veins run in the roof of the 3rd ventricle backwards to form **the great cerebral vein**, which drains into **the straight sinus.**
With addition the inferior sagittal sinus




- 1- Deep middle cerebral vein.
- 2- Anterior cerebral vein.
- 3- Striate vein.
- B.: Basal vein.
- I.: Internal cerebral vein.
- G.: Great cerebral vein.

6. Great cerebral vein (vein of Galen):

- Is the main vein of the medial surface of C.H.
- Formed by union of **the two internal cerebral veins.**
- **The great cerebral vein** unites with **the inferior sagittal sinus** to form **the straight sinus.**
- It receives also **the two basal veins.**





The Cerebral Dural Sinuses

- The dural venous sinuses (also called dural sinuses, cerebral sinuses, or cranial sinuses) are venous channels found between two layers of dura mater in the brain.
- They receive blood from superficial and deep veins of the brain.
- They receive CSF from the subarachnoid space (by the arachnoid granulation and villi).
- The dural sinuses finally connected with the internal jugular vein. Then to brachiocephalic vein then right atrium.
- There are at least three single and six paired dural sinuses.


So if occlusion happens it may cause bilateral symptoms and they came with hemorrhage and bleeding. Cuz there is no drains to the cerebral veins so it will increase the venous pressure and rupture all the veins. The treatment is full dose coagulant to open the sinus
على الرغم من انه في ترويق

As they are single so they would be in the middle.

The Single Venous Sinuses


1. **The superior sagittal sinus:** usually terminates in **the right transverse sinus.**
2. **The inferior sagittal sinus:** joins the great cerebral vein to form **the straight sinus.** With great cerebral vein.
3. **The straight sinus “tentorial or rectus sinus”:** formed by the union of **the great cerebral vein and the inferior sagittal sinus,** and usually drains into **the left transverse sinus.** Drains to sigmoid Then to internal jugular vein.

The Dura Mater and Dural Sinuses




(a) Midsagittal view

Absorption the CSF from
subarachnoid space to dural sinuses




Meninges and Diploic Veins


Coronal Dissection



The Paired Sinuses Rt and Lt

1. **The sphenoparietal sinus:** drains into **the cavernous sinus**.
2. **The cavernous sinuses:** drains into **the superior and inferior sinuses**.
3. **The superior petrosal sinuses:** extends from the cavernous sinus to the junction of **the transverse and sigmoid sinus**.
4. **The inferior petrosal sinuses:** drains the cavernous sinus **into the sigmoid sinus**.
5. **The transverse (lateral) sinuses:** drains into **the sigmoid sinus**.
6. **The sigmoid sinuses:** a continuation of the transverse sinus at the superior petrosal sinuses and drains into **the internal jugular vein**.





The Cavernous Sinuses


□ Contains within the **lateral wall** of the cavernous sinus:

1. The oculomotor nerve.
2. The trochlear nerve.
3. The ^{V1}ophthalmic division of the trigeminal nerve.
4. The ^{V2}maxillary division of the trigeminal nerve.

□ Contains within the **medial wall** of the cavernous sinus:


1. The abducent nerve.
2. The internal carotid artery.

Cavernous sinus thrombosis → painful ophthalmoplegia.




Cavernous Sinus

Coronal Section



The Cerebral Dural Sinuses

Name	Drains to
Inferior sagittal sinus	Straight sinus
Superior sagittal sinus	Typically becomes right transverse sinus or confluence of sinuses
Straight sinus	Typically becomes left transverse sinus or confluence of sinuses
Occipital sinus	Confluence of sinuses
Confluence of sinuses	Right and Left transverse sinuses
Sphenoparietal sinuses	Cavernous sinuses
Cavernous sinuses	Superior and inferior petrosal sinuses
Superior petrosal sinus	Transverse sinuses
Transverse sinuses	Sigmoid sinus
Inferior petrosal sinus	Sigmoid sinus
Sigmoid sinuses	Internal jugular vein




The **confluence of sinuses** is the connecting point of the superior sagittal, and straight sinuses.

Classification of Stroke


❖ *Occlusion of arteries* **Ischemic Stroke: 85%.**

- Thrombotic.
- Embolic. *Migratory thrombus*
- Lacunar.
- Hemodynamic. *watershid syndrome*

Thrombotic Stroke




Embolic Stroke






❖ *Rupture of blood vessels.* **Hemorrhagic Stroke: 15 %.**

- Intracerebral and intracerebellar hemorrhage 10%.
- Subarachnoid hemorrhage 5%.
in subarachnoid space.

Cerebral Hemorrhage







Rupture of large artery cause subarachnoid hemorrhage
 If penetrating artery like charcots artery it will cause intracerebral hematoma

Occlusion Internal Carotid Artery ICA

1. The symptoms and signs are those of **MCA** including contralateral hemiparesis and hemianesthesia.
2. There is **partial** or **complete loss of vision** on the same side, but permanent loss is **rare** (emboli dislodged from the internal carotid artery reach the retina through the ophthalmic artery).
الشبكية Gives retinal artery.

Occlusion of the main Anterior Cerebral Artery

- Contralateral weakness and hyperreflexia in the lower extremity.
- Contralateral somatosensory loss in the lower extremity with paresthesia, numbness, and loss of vibration sensation.
- Urinary incontinence with bilateral infarction.
- Apraxia. Loss of motor skills
- Mentality and personality changes.

Occlusion of the main Middle Cerebral Artery MCA

- Contralateral facial weakness and weakness and loss of sensation in the upper extremity and face “**Contralateral faciobrachial type hemiplegia and hemihypoesthesia**”.
- Loss of sensory discrimination and stereognosis.
- **Hemineglect** (may occur with either left or right parietal lesions).
- **Conjugate deviation** of the eyes to the affected side
- Frontal lobe syndrome.
- **Brocca's** expressive aphasia, agraphia. Nonfluent
- **Wernicke** “sensory” aphasia. Fluent
- Contralateral homonymous hemianopia) damage to the optic radiation. (فقدان نصف مجال الابصار)
- Anosognosia. المريض متجاهل المرض وغير واعى انه مريض


Occlusion of the main Posterior Cerebral Artery PCA

- **Unilateral Lesion of occipital lobe:** **contralateral homonymous hemianopia** with **macular sparing** due to the occipital pole receiving Macular area take dual blood supply collateral blood supply from the middle cerebral artery.
- **Bilateral Lesion of occipital lobe:** **cortical blindness** (pupils are reactive to light).
- **Irritation of occipital lobe:** Visual hallucination.
- Memory deficit (**amnesia**).
- Visual agnosia.
- **Thalamic syndrome.** Contralateral pain hemi type
Contralateral anesthesia
Emotional liability
Decrease consciousness

Vertebrobasilar Artery Occlusion

Damage of trigeminal sensory nucleus

1. Ipsilateral pain and temperature sensory loss of the face and contralateral pain and temperature sensory loss of the body
2. Attacks of hemianopia or complete cortical blindness.
3. Bulbar syndrome. *صعوبة في البلع والكلام*
4. Vertigo, nystagmus, nausea, and vomiting
5. Ipsilateral Horner syndrome
6. Ipsilateral ataxia and other cerebellar signs
7. Unilateral or bilateral hemiparesis
8. Coma



The Cerebro-Spinal Fluid

Definition: it is the fluid which circulates in the ventricles and central canal of the CNS and also fills the subarachnoid space.

Production of the C.S.F.:


- The C.S.F. is secreted by the ependymal cells covering the choroid plexuses of the ventricles, mainly the lateral ventricles.
- The ependymal cells extract the CSF from the capillaries and secrete it into the cavity of the ventricles.


Formation of Cerebrospinal Fluid

- **CSF is formed mainly** in the choroid plexuses of the lateral, 3rd & 4th ventricles.
- Some originates from **the ependymal cells** lining the ventricles & from the brain substance through the perivascular spaces.
- The choroid plexuses actively secrete CSF.

Circulation of the Cerebro-Spinal Fluid


The CSF in each lateral ventricle → interventricular foramen of
Monro → third ventricle → aqueduct of Sylvius → fourth
ventricle → some fluid enter the central canal of spinal cord but
the majority passes through the 3 foramina (one median of
foramen Magendie to cerebellomedullary cistern of
subarachnoid space & two lateral foramina of Luschka to
pontine cistern of subarachnoid space) → subarachnoid space
where it flows over the surface of the brain and spinal cord.





Absorption of the Cerebro-Spinal Fluid

- The CSF circulates in the subarachnoid space.
- Absorbed by **arachnoid villi and granulations** which absorbs the C.S.F. from the subarachnoid space into the dural venous sinuses, **especially the superior sagittal sinus** to reach the blood stream.



Functions of the Cerebro-Spinal Fluid

1. Support and protects the CNS against trauma.
2. It takes the place of the lymph in the CNS which is devoid of lymphatic vessels.
3. Regulation of the intracranial pressure.
4. Removes metabolites and toxics from the central nervous system.
5. Nourishes the central nervous system.

Characters of the Normal Cerebro-Spinal Fluid

- **Appearance:** clear watery fluid, isotonic.
- **Amount:** 100- 150 ml.
- **Replacement:** it continuously produced and absorbed at a rate of 400-500 ml daily (it is replaced 3 times per day).
- **Pressure:** 100-180 mm of water.
- **Rate of production:** 0,3-0,4 ml/min.
- **Cells:** 0-5 WBCs per cubic millimeter.
- **Glucose:** 60 mg/dl (70% of serum glucose).
- **Total protein:** 15-45 mg/dl.

Formation of CSF by the Choroid Plexus Epithelium



- Lipid-soluble substances (CO_2 and O_2) and H_2O freely cross the blood–brain barrier and equilibrate between blood and CSF.
- **Protein and cholesterol** are **excluded** from the CSF because of their **large molecular size**.
- Other substances are transported by carriers in the choroid plexus epithelium.
- They may be secreted from blood into the CSF or absorbed from the CSF into blood.

CSF Sampling

- **To analyse the constituents of the CSF:**
 - Meningitis
 - Malignant disease
 - Multiple sclerosis and other neurological diseases
- **To remove CSF to reduce intracranial pressure:**
 - For Surgical access
 - Pseudotumor cerebri


Hydrocephalus

- **Definition:** Distension of the ventricular system (Ventriculomegaly) due to inadequate passage of CSF from its point of production to the point of absorption”.
- **3 causes:**
 1. Choroid plexus papilloma: Rarely over-secretion of CSF.
 2. Can be from Obstruction to flow of CSF from the ventricles into the subarachnoid Space (**Obstructive**)
 - Masses in ventricles or adjacent brain.
 - Developmental membranes or stenosis of aqueduct.
 3. Can be from failure of absorption (**Communicating**)
 - Failure of the absorbing mechanism usually after a change in CSF contents e.g. acutely after Meningitis or SAH, Chronically also. Cryptogenic, congenital



Raised CSF Pressure & Papilledema

- The optic nerves are surrounded by sheaths derived from the pia mater, arachnoid mater, and dura mater.
- A rise of CSF pressure will compress the thin walls of the retinal vein.
- This will result in congestion of the retinal vein, bulging forward of the optic disc, and edema of the optic disc (papilledema).
- Persistent **papilledema** leads to **optic atrophy and blindness**.



Thank You
Thank You

