

Khalleedd✓

Lecture

Main Functions of the Cranial and Spinal Nerves

Dr. Eyad M. Hussein

**Ph.D. of Neurology
Consultant & Head of Neurology Department,
Nasser Hospital**

The Cranial Nerves

Efferent carry motor information away from CNS represents by Brain
While Afferent carry information found all over the body toward the CNS

- The cranial nerves are consist of 12 pairs of nerves that arise from the brain.

The difference between spinal and cranial nerves is:

1. The 1st enter or exit the skull through intervertebral foreman while the 2nd enters or exit through specialized foremen in the skull
2. All of the spinal nerves are mixed contains motor sensory and may have autonomic while the cranial nerves divided to 3 groups pure motor, pure sensory, and mixed.

- They ^{Motor}exit/^{Sensory}enter the cranium through openings in the skull.
_{Efferent Afferent}

- The olfactory, optic and vestibulocochlear nerves are purely sensory nerves “special sensory” (128).

- The oculomotor³, trochlear⁴, abducent⁶, accessory¹¹ and hypoglossal¹² nerves are purely motor nerves. 3-4-6-11-12

- The trigeminal, facial, glossopharyngeal and vagus nerves are mixed nerves (10975).

Cranial nerves contain autonomic fibers 10973

سؤال بيطرح نفسه انوال ١٠ وال ٩ وال ٧ حطيناهم مع ال mixed ليسوا حطينت نيف رقم ٣ مع ال mixed
الجواب بلك سهولة انوال motor
↓

Supplies constrictor pupil muscle and Ciliary muscle that play a role in accommodation reflex.

Functional Modalities

- voluntary • **Somatic:** skeletal muscle, joint, tendon, ligament, and skin.
- involuntary • **Visceral:** smooth muscles, cardiac muscles, and glands. GIT peristalsis.
 - **Sensory “Afferent”:**
 - ✓ **General.** Pain sensation
 - ✓ **Special.** Taste sensation
 - **Motor “Efferent”:**
 - ✓ **General.**
 - ✓ **Special.**
- **Organic sensations:** e.g. hunger, thirst and sexual sensations **“Hypothalamic Sensations”.**
The center is found in it and the receptors

Motor Functional Modalities

1. General Somatic Efferent (GSE): somatomotor, or

Found in anterior somatic horn of spinal cord, give the signal via spinal nerve and supply skeletal muscles

somatic motor fibers, arise from motor neuron cell bodies in the AHCs of the spinal cord, which supply the skeletal muscles.

2. General Visceral Efferent (GVE): visceral or autonomic

Involuntary motor

efferent, are the efferent nerve fibers of the autonomic nervous system (motor innervation to smooth muscle, cardiac muscle, and glands). For internal organs

3. The Special Visceral Efferent (SVE): which supply

Do not supply smooth muscles

muscles of the trigeminal, facial, glossopharyngeal, vagus, and cranial accessory nerves.

Supplies somatic skeletal muscles and follow the cranial nerves

special visceral efferents, which innervate striped muscles of branchial origin, and general visceral efferents, which innervate involuntary muscles and secreting glands

Sensory Functional Modalities

Afferent

1. **General Somatic Afferent (GSA): somatic sensory "afferent"**
For touch pain proprioception temperature sensation
fibers arise from neurons in sensory ganglia and are found in all the spinal nerves, and conduct general sensations from the surface of the body through the dorsal roots to the spinal cord.
2. **Special Somatic Afferent (SSA): afferent nerves of the special senses,** (fibers include the optic nerve and the vestibulocochlear nerve).
For special sensation not related to GIT = hearing and vision
Vision Hearing
3. **General Visceral Afferent (GVA): conduct sensory impulses (usually pain or reflex sensations)** from the internal organs, glands, and blood vessels to the central nervous system.
For sensory impulses of internal organs
Like the stretched urinary bladder sensation Irritation from diseased or inflamed organ cause pain sensation in it
4. **Special Visceral Afferent fibers (SVA): afferent fiber that develop in association with the gastrointestinal tract.** They carry the special senses of smell (olfaction) and taste (gustation). The cranial nerves containing SVA fibers are the olfactory nerve (I), the facial nerve (VII), the glossopharyngeal nerve (IX), and the vagus nerve (X).
For special sensation related to GIT = smell and taste
Smell Taste

Taste sensation by 3 cranial nerves: *Facial nerve by chorda tympani for ant. 2/3 * Lingual of glossopharyngeal for post. 1/3 * Vagus give superior laryngeal and has external and internal branches the internal supplies the sensation above vocal cords and give a branch for taste sensation in epiglottis.

The Main Three Functions of the Cranial Nerves

1. **General Motor & Sensory Innervation:** of the skin, muscles, and joints in the head and neck region.

2. **Special Senses:** ^{Somatic} vision, hearing, ^{visceral} smell, & Taste.

3. **Autonomic “Visceral” Functions”:** via

parasympathetic innervation to the visceral organs.

^{Salivation}
Parotid gland works to secrete saliva: by tympanic nerve tympanic nerve from glossopharyngeal before it reach the gland it relay in otic ganglion
Sublingual and submandibular salivary glands works by Corda tympani nerve before that it relay in submandibular ganglion
How the ^{tear} mechanism works by GSPN greater superficial petrosal nerve before that it relay in sphenopalatine ganglion
↓
Branch of facial nerve

Classification of the Cranial Nerves

No	Name	Function	Site of nucleus	Opening in the skull
1	Olfactory	Smell	Temporal Lobe	Cribriform plate
2	Optic	Vision	Occipital lobe	Optic canal
3	Oculomotor	Motor	Midbrain	Superior orbital fissure
4	Trochlear	Motor	Midbrain	Superior orbital fissure
5	Trigeminal: Ophthalmic ^{v1} Maxillary ^{v2} Mandibular ^{v3}	Sensory Sensory Mixed	Midbrain, Pons Pons Medulla oblongata Upper C. spinal cord	Superior orbital fissure Foramen rotundum Foramen ovale
6	Abducent	Motor	Pons	Superior orbital fissure
7	Facial	Mixed	Pons	Internal acoustic meatus, Facial canal, Stylomastoid foramen
8	Vestibulo- cochlear	Hearing and balance	Pons	Internal acoustic meatus
9	Glossopharyngeal	Mixed	Medulla oblongata	Jugular foramen
10	Vagus	Mixed	Medulla oblongata	Jugular foramen
11	Accessory	Motor	Medulla oblongata C1-C4 AHC of SC	Jugular foramen
12	Hypoglossal	Motor	Medulla oblongata	Hypoglossal canal ⁸

Cranial Nerves

Motor → 3-4-6-11-12

Sensory → 1-2-8

Mixed → 10-9-7-5

Autonomic → 10-9-7-3

With exception (3) is pure motor even the autonomic fibers

Cerebral hemispheres → 2 → temporal (1) occipital (2)

Midbrain → 2 → (3) + (4)

Pons → 4 → (5) + (6) + (7) ((8)

Medulla oblongata → 4 → (9) + (10) + (11) + (12)


With some exceptions.

Parts of trigeminal (5) nerve nucleus is found in midbrain

And parts of them descends to the spinal cord.

Parts of facial (7) nerve nucleus is found in the medulla oblongata.

Accessory nerve (11) has cranial and spinal parts the 2nd is in the anterior horn of the upper cervical segment.



They arranged according to the enter or the exit from the brain

Functions:

* Sensory

1 → smell

2 → vision

8 → hearing and pulse

* mixed

5

7

9

10

* motor

3

4

6

11

12

Opening in the skull

Olfactory → cribriform plate may arise from it olfactory groove meningioma and cause Foster-Kinney syndrome.

Optic → optic canal.

Oculomotor, Trochlear, Ophthalmic branch of Trigeminal, and Abducent → superior orbital fissure.

Maxillary → foramen rotundum

Mandibular → foramen ovale

Facial → enter through internal acoustics meatus and exit through stylomastoid foramen.


Vestibulo-cochlear → internal acoustics meatus.

Glossopharyngeal, vagus, accessory → jugular foramen

Hypoglossal → specific canal

Brainstem

Anterioinferior View



Notes on the slides before

Oculomotor ventral surface anterior of mid brain

Trochlear dorsal surface posterior of midbrain

Trigeminal has large root sensory and small root autonomic

Facial has two roots large is motor and small sensory and parasympathetic

In cerebellum pontine angle pass 3 cranial nerves 5th 7th and 8th if damaged called CPA syndrome
اجت للدكتور حالة حكاها وكانت بتشكي من اشياء يعملوها هقول ال 3 نيفز وبلش الكومبلايه بفقدان السمع وزنه في الداء وبعدها بلسن اعراض النيفيه الثانيه، والدكتور شخصها انها ورم في
العصب التامع وعامله ضغط عال السابع والخامس

Accessory has 2 roots cranium part from medulla oblongata and spinal parts from spinal cord and supplied trapezius muscle and sternocleidomastoid muscle

I. The Olfactory Nerve

Its not a single tract it composed of 20 filaments on each side and enters through cribriform plate on alfactory groove and deliver the signal to olfactory pulp then olfactory tract and attached to the inf. Surface of cerebellar hemisphere

The pathway

- **Olfactory Nerve:** (Latin for "to smell").
- **Function and modality:** **Special visceral afferent** (smell sensation).
Means sensory
Related to git
- **Olfactory tract** located in the anterior cranial fossa and attached to the inferior surface of frontal lobe.
So the signal reach the 2 hemispheres
The right goes to Rt and Lt and vice verse




II. The Optic Nerve

❖ Optic Nerve: (Latin for "to see").

- A bundle of 1 million nerve fibers.
- Responsible for transmitting nerve signals from the retina to the brain. Occipital lobe for visual center
- The optic disc is the front surface of the optic nerve.


❖ **Function and modality:** special somatic afferent (sense of vision) and pupillary light reflex.




Pathway of the Vision

Receptors of light are **rods and cones of retina** → **optic nerve** → optic canal of the sphenoid bone → **optic chiasma** (the medial fibers decussate to the opposite optic tract, while the lateral fibers continue in the same optic tract) → **optic tract** (gives fibers to **Edinger-Westphale nucleus in midbrain for light reflex**) → relay in the lateral geniculate body → posterior limb of internal capsule → optic radiation → end in areas **17, 18, 19 of occipital lobes.**

Visual pathway field defects






Light reflex

Direct light reflex if we put a light in the right eye it will cause constriction to the same pupil that exposed to light

Indirect light reflex if we expose right eye to a light it will cause the left eye pupil to constricts

Pathway of vision and light reflex




Sensory

Motor

The nerves that responsible for it is the optic as Afferent and oculomotor as Efferent

The Visual Pathway




III. The oculomotor Nerve

E in efferent means motor

- **Oculomotor nerve:** (*Latin for "eye" and "moving"*).
- **Function and modality:**
 1. **General somatic efferent:** supplies extraocular ^{voluntary} “skeletal muscles” for the eye ball movement.
 2. **General visceral efferent:** supplies intraocular smooth muscles for pupillary reaction and light reflex.
Sphincter pupil muscle + ciliary muscle
- **Site of nucleus:** it lies in the tegmentum of midbrain.
- Emerges from the ventral ^{Anterior} surface of the midbrain.
- **Foramen:** superior orbital fissure **“SOF”** of the sphenoid bone.

It pass between posterior cerebral artery and superior cerebella artery is aneurysm happened and it cause pressure on the nerve it will cause oculomotor nerve palsy



The oculomotor N. supplies all of the intra- and extraocular muscles, Except:


حناخذهم كماد شوية شو بيعملو ويكيف بيحركو العييه ف اعرفو هاد بس انهم ما يتغذوا بالنيرف رقم ٣ وخلص

1. **Superior oblique muscle** → from the trochlear nerve – IV.
2. **Lateral rectus muscle** → from the abducent nerve – VI.
3. **Dilator pupillae muscle** → from the sympathetic fibers.
4. **Muller's muscle** → from the sympathetic fibers.

If sympathetic is damaged it will cause partial ptosis with Horner syndrome patients

Horner → ptosis, miosis, and anhidrosis

But if oculomotor is damaged it will cause complete ptosis because it supply the main muscle that responsible for.



Structure of Oculomotor Nuclei

1. **The Edinger-Westphal nucleus:** parasympathetic function which supplies two intraocular muscles:


- a. **Constrictor pupillae muscle** → Miosis. *Constriction of the pupil.*
- b. **Ciliary muscle** → Responsible for light and accommodation reflexes.

2. **The lateral cell mass (motor function):** supplies five extra ocular muscles:

- a. **Levator palpebrae superioris muscle.** *Elevate the eyelid and open the eye*
- b. **Superior rectus muscle.**
- c. **Medial rectus muscle.** *There is lateral rectus muscle supplied by abductors*
- d. **Inferior oblique muscle.** *There is superior oblique muscle supplied by trochlear*
- e. **Inferior rectus muscle.**

3. **The central nucleus of Perlia:** motor function which supplies **the medial recti muscles of the two eyes** allowing to contract together when both eyes converge to look to a near object.

Plus accommodation reflex by ciliary muscle



Branches of the Oculomotor Nerve

1. Superior division “Somatic motor and sympathetic”:

which supplies the:

- a. Levator palpebrae superioris muscle.
- b. Superior rectus muscle.

2. Inferior division “Somatic motor and parasympathetic”:

which supplies the:

a. Medial rectus muscle.



b. Inferior rectus muscle.




c. Inferior oblique muscle.



d. Constrictor pupillae muscle (via ciliary ganglion).


Sphincter pupil muscles

e. Ciliary muscle (via ciliary ganglion).



Nerves of Orbit - Muscles Partially Cut Away

Superior View



IV. The Trochlear Nerve

Arise from a dorsal surface (Posterior) of midbrain.

- It is purely a motor nerve and the smallest cranial nerve.
- **Function and modality:** ^{Motor} general somatic efferent → supplies the superior oblique muscle, which abducts, depresses and internally rotates the eye
- **Site of the nucleus:** it lies in the tegmentum of the midbrain.
- Emerges from the posterior surface of the midbrain.
- **Foramen:** superior orbital fissure “SOF” of the sphenoid bone.


Oculomotor and trochlear enters to the upper lateral wall cavernous sinus then to superior orbital fissure and supply sup. Oblique muscle. While abducent nerve in the upper medial wall cavernous sinus.


VI. The Abducent Nerve

- **Abducent nerve:** *(Latin for "abduction")*.
- **Function and modality:** **general somatic efferent** ^{Motor} → supplies **the lateral rectus muscle**.
Then it pass in the medial wall of cavernous sinus then goes to superior orbital fissure and supply lateral rectus muscle → for abduction to eye.
- **Site of the nucleus:** it lies in pons. ventral surface (anterior)
- The abducent nerve emerges from the lower border of the pons (between the pons and medulla oblongata).
- **Foramen:** superior orbital fissure **“SOF”** of the sphenoid bone.

Brainstem


Anterioinferior View





The Actions of the Ocular Muscles

كل العضلات يتمشي ع هوى الاسم ما عدا الي الها علاقة بال oblique



Rectus muscle

Superior → elevation

Inferior → depression


Lateral → abduction


Medial → adduction

1. **Lateral Rectus Muscle:** abducts (laterally) the eye ball.
2. **Medial Rectus Muscle:** adducts (medially) the eye ball.
3. **Superior Rectus Muscle:** elevates, **adducts** and rotates medially.
4. **Inferior Rectus Muscle:** depresses, **adducts** and rotates medially.
5. **Superior Oblique Muscle:** depresses, **abducts** and rotates laterally. *It may do adduction*
6. **Inferior Oblique Muscle:** elevates, **abducts** and rotates laterally.


Extrinsic Eye Muscles


Right Lateral View






Right Eye, Anterior View






superior rectus
Elevation with adduction




inferior oblique
Elevation with abduction




inferior rectus
Depression with adduction



superior oblique
Depression with abduction



Adduction
medial rectus



Abduction
lateral rectus

Lesion of the Oculomotor Nerve

Main muscle for elevation of the eye and open it if paralyzed it will cause ptosis

- **Ptosis:** paresis of the levator palpebrae superioris muscle.
- **Diplopia:** occurs only on elevation of eye lid.
Double vision
- **Squint:** divergent paralytic.
انحراف بالعين
- **Mydriasis:** dilated fixed pupil.
- **Loss of light and accommodation reflexes.**

cause of lateral rectus work it will cause abduction of the eye and medial rectus doesn't work so it will pull the eye laterally
if there is ptosis it will not cause diplopia




الحالة المرضية التي عندهم diplopia التي يسببها تضادة واسترخاء العينين
وحدة وبالتناوب كل يوم عينه، عشاء انخلص منه ال double vision

Because the constriction that caused by parasympathetic is deleted so the sympathetic is dominant so cause dilated pupil.



Lesion of the Oculomotor Nerve



If a patient came with oculomotor nerve palsy either she is DM with mono neuropathy or aneurysm compression to the nerve

If case presentation with one question and pupil is intact it is believed that it is for DM that affect the axons and autonomic fibers is more external but if he has mydriasis it is believed to be compression or pressure to the nerve cause its the 1st symptom appears

According to the place of the compression:

If the problem in cavernous sinus where ophthalmic and maxillary passes it will cause painful ophthalmoplegia if it was by superior orbital fissure tumor it will cause the same because of the ophthalmic branch of trigeminal nerve

But if the nerve is just affected it will be painless

Lesion of the Trochlear Nerve

1. Diplopia on looking down and out.
2. Limitation of movement during examination for eye movement down and out. Depression and abduction

Lesion of the Abducent Nerve

1. Diplopia on looking out wards.
2. Limitation during examination of eye movement in the outward direction. Abduction
3. Convergent paralytic squint.



Light Reflex Done ✓

- **Direct Light Reflex:** Light in one eye leads to constriction (miosis) of the **same** eye
- **Indirect Light Reflex :** Light in one eye leads to constriction (miosis) of the **opposite** eye.

Pathway of Light Reflex

Fibers from rods and cones of retina → optic nerve → optic chiasma → optic tract → before reaching the lateral geniculate body → they pass to the third cranial (oculomotor) nerve nucleus (Edinger-Westphal nucleus) in midbrain of both side → through oculomotor nerve → ciliary ganglion → supply the constrictor pupillae muscle

Parasympathetic




Accommodation and Convergence Reflex

Done ✓

Patient is asked to follow your finger from a far point to a near one:

- ❖ Convergence (contraction of both medial recti muscle).
- ❖ Miosis (contraction of constrictor pupillae muscle).

ciliary muscle




V. The Trigeminal Nerve

- The trigeminal nerve is **mixed nerve** (motor and sensory fibers).
- The trigeminal nerve is more important as a sensory nerve. **It carries:**
 - a. Pain, touch, and temperature sensations.
 - b. Proprioception (Deep) sensation.
- It is the thickest cranial nerve and arises by two roots:
 - a. **Large Root:** sensory function. Pass through cerebello pontine angle with 7th and 8th
 - b. **Small Motor Root:** motor function.
- The two roots are attached to the **ventro-lateral aspect** of the pons. Anterolateral


Brainstem

Anterioinferior View



Nuclei of the Trigeminal Nerve

1. **Main Sensory Nucleus:** lies in the middle part of the pons (for touch sensation).
2. ^{Long nucleus} **Spinal (Descending) Nucleus:** lies in the medulla oblongata and extends upwards to the lower part of the pons and downwards to the upper cervical spinal cord (for pain and temperature sensation).
3. **Mesencephalic Nucleus:** lies in the ^{Exception} midbrain and upper part of the pons (deep sensation).
4. **Motor Nucleus (motor function):** lies in the middle of the pons. It give motor fibers that joins with the mandibular nerve.
*For 8 muscles
4 mastication and other 4 non mastication.*
The only one that has motor function in 5th cranial nerve.



Branches of the Trigeminal Nerve

The trigeminal nerve has three branches:

1. Ophthalmic Nerve (V1): **sensory.**

2. Maxillary Nerve (V2): **sensory.**


3. Mandibular Nerve (V3): **mixed.**

Supplies 8 muscles

4 mastication

4 non mastication

Ophthalmic and Maxillary Nerves



The Ophthalmic Nerve (V₁)

- It is the first branch of the trigeminal nerve.
- **Function and modality:** general somatic ^{Sensory} afferent.
- **Foramen:** superior orbital fissure “SOF” of the sphenoid bone.
- **It supplies the:**
 1. Skin over forehead.
 2. Skin over upper eyelid.
 3. Eyeball, conjunctiva, & lacrimal gland.
 4. Corneal reflex.
called conjunctiva reflex
*لو جنبنا جسم غريب على سبيل المثال قطنة ومزناها على سطح العينه
حتلاقي انو عينك سكرت لحالها وهادي هيا ال corneal reflex*
 5. Nasal mucosa.
 6. Frontal sinus.
 7. External nose.
 8. Dura mater in the anterior cranial fossa.

2 cranial nerves are responsible
sensory V1 and V2 while motor facial nerve 7th
Afferent Efferent

The Maxillary Nerve (V₂)

- It is the second branch of the trigeminal nerve.
- **Function and modality:** general somatic ^{Sensory} afferent.
- **Foramen:** rotundum foramen.
- **It supplies the:**

- Maxillary area of skin of the face.
- Maxillary teeth (of the upper jaw).
- Mucosa membrane related to maxilla.
- Maxillary sinus.

Irritation of soft palat and cause elevation of it and cause involuntary swallowing

- Palatal reflex.

Sensory → maxillary V₂ and glossopharyngeal 9th
Motor → glossopharyngeal 9th and vagal 10th

- Dura matter in the middle cranial fossa.

The Mandibular Nerve (V₃)

- It is the third branch of the trigeminal nerve.
- **Functions and modalities:** mixed nerve **general somatic**
Sensory **afferent** & **special visceral** Motor **efferent**.
- **Foramen:** rotundum foramen.
- **Origin:** They joined each other it is a mixed nerve formed of two roots:
 1. **Sensory root:** arises from trigeminal ganglion, runs forwards and laterally to the **foramen ovale**.
 2. **Motor root:** a smaller root which arises from trigeminal motor nucleus in the pons. It passes deep to the trigeminal ganglion to join the sensory root in the **foramen ovale**.

Sensory Functions of the Mandibular Nerve


❖ The sensory part of the mandibular nerve carries general sensory information from the:

1. Mucous membranes of the mouth and cheek.
2. Anterior two-thirds of the tongue.
3. Lower teeth.
4. Skin of the lower jaw.
5. Jaw jerk.
6. Side of the head and scalp.
7. Meninges of the middle cranial fossa.

Motor Functions of the Mandibular Nerve

1. Motor fibers to muscles of mastication:

- Temporalis muscle.
- Masseter muscle.
- Medial pterygoids muscle.
- Lateral pterygoids muscle.




2. Motor fibers to other four muscles:

- Mylohyoid muscle.
- Anterior belly of digastric muscle.
- Tensor palati muscle (from otic ganglion).
- Tensor tympani muscle (from otic ganglion).

The posterior branch supplied by a branch of facial nerve

Dermatomes of Head and Neck



From ophthalmic division
of trigeminal nerve (V1)

Supraorbital nerve

Supratrochlear nerve

Palpebral branch
of lacrimal nerve

Infratrochlear nerve

External nasal branch of
anterior ethmoidal
nerve

From maxillary division
of trigeminal nerve (V2)

Infraorbital nerve

Zygomaticofacial nerve

Zygomaticotemporal nerve

From mandibular division
of trigeminal nerve (V3)

Mental nerve

Buccal nerve

Auriculotemporal nerve

Auricular branch
of vagus nerve (X)

Medial branches
of dorsal rami
of cervical spinal
nerves

Greater occipital
nerve (C2)

3rd occipital
nerve (C3)

From 4th, 5th,
6th and 7th
nerves in
succession below


Branches from
cervical plexus

Lesser occipital
nerve (C2, 3)

Great auricular
nerve (C2, 3)


Transverse cervical
nerve (C2, 3)

Supraclavicular
nerves (C3, 4)



Trigeminal Neuralgia (TN)

“Tic Douloureux”



- TN is defined as paroxysmal, ^{كلا المبرتي بييجو ينفصل ال complain} stereotyped attacks of severe sharp pain (**stabbing or electric-shock like pain**) in the distribution of one or more branches of the trigeminal nerve.

VII. The Facial Nerve

- It is a mixed nerve which contains motor, sensory and parasympathetic fibers.
- It is more important as a motor than as a sensory nerve.
- **Modalities:** GVE, SVE, GSA, & SVA.
- **Foramen:** Internal auditory meatus → Stylomastoid foramen.

Motor part

Sensory part

① UMNL upper motor neuron lesion → C.B.T damage

② LMNL lower motor neuron lesion → nuclear or nerve lesion

① if damaged result contralateral lower part paralysis

② if damaged result ipsilateral hemifacial paralysis

Roots of the Facial Nerve

➤ **Facial Nerve (motor nerve):** is a purely motor nerve supplies all the muscles of face expression.


➤ **Nerve Intermedius:** consists of sensory and parasympathetic fibers.

In general the C.B.T gives dorsal and ventral contralateral in addition to dorsal ipsilateral

Dorsal part cause its importance for the eye has dual innervation from both cerebral hemispheres.


D → dorsal (anterior) part of nucleus for upper part of face

V → ventral (posterior) part of nucleus for lower part of face



Brainstem

Anterioinferior View



Nuclei of the Facial Nerve

1. **Facial motor nucleus (Motor)**: lies in **the pons** and supplies the facial muscles.
2. **Superior salivary nucleus (Parasympathetic)**: lies in **the pons** for:
 - All salivary glands except parotid gland supplied by glossopharyngeal 9th*
 - a. Submandibular and sublingual salivary glands (**Chorda tympani nerve**). *Branch of facial*
 - b. Glands of the palate, nose, lacrimal glands and nasopharynx (**Greater superficial petrosal nerve**).
3. **Solitary nucleus**: lies in **the medulla oblongata** for:

7th	◦ General sensation from the skin of the external auricle and tympanic membrane.
9th	
10th	◦ Taste sensations from the anterior 2/3 of the tongue.

Functions of the Facial Nerve

1. Motor function to muscles of facial expression.
2. Parasympathetic to all glands of head (submandibular, sublingual and lacrimal glands) except the parotid gland (glossopharyngeal nerve).
3. Sensory for skin of the external auricle and tympanic membrane.
4. Taste for anterior 2/3 of the tongue.
5. Corneal reflex.
called conjunctiva reflex

لو جينا جسم غريب على سبيل المثال قطنة ومزناها على سطح العين
حتلاقي اناو عيّنك سكرن لخالها وهادي هيّا ال corneal reflex

2 cranial nerves are responsible
sensory V1 and V2 while motor facial nerve 7th
Afferent Efferent

Branches of the Facial Nerve

Inside the Fallopian Canal

10 branches

3 intracanal

2 immediately after exit

5 final branches


1. The greater superficial petrosal nerve “GSPN”.
Only parasympathetic

2. Small branch to the ^{على اسم العظمة المتصلة فيها}stapedius muscle.
Only motor
In internal ear its contraction decrease transmission of the high vibrations and sound from external ear to internal

3. The chorda tympani.
Mixed: sensory and parasympathetic

For taste sensation of ant. 2/3 of the tongue

And parasympathetic fibers for submandibular ganglion for submandibular and sublingual salivary gland.



Its outlet of the skull.

Branches of the Facial Nerve

After it Leaves the Skull

❖ Immediately after it leaves the skull “2 branches”:

1. The posterior auricular nerve. Carry sensation of the auricle and supply occipital belly of occipitofrontalis muscle
2. The nerve to the posterior belly of the digastric muscle and to the stylohyoid muscle.

The nerve reach's posterior medial surface of parotid gland

❖ Terminal branches of the facial nerve “5 branches”:


1. Temporal nerve (from temporofacial branch).
2. Zygomatic nerve (from temporofacial branch).
3. Buccal nerve (from cervicofacial branch). Supply 2 muscles: orbicularis oris and buccinator
4. Marginal mandibular nerve (from cervicofacial branch). Supply muscle around the mandibular angle.
5. Cervical nerve (from cervicofacial branch). Supply laryngeal muscle


Upper part for orbicularis muscle for its importance has dual innervation

Upper and lower

Upper and lower
orbicularis oculi muscle
Close the eye

For nasal muscle and muscle
between eye and mouth





Bell's Palsy

- ❖ **Sir Charles Bell “Scottish Surgeon”**
first described the anatomy and function
of the facial nerve in the 1800.
- ❖ **Bell's palsy:** defined as an acute
peripheral facial nerve palsy of
unknown cause.
- ❖ The annual incidence rate is between
13-34 cases per 100,000 population.
- ❖ **Etiology:** Herpes simplex virus
reactivation.



(12 November 1774 – 28 April 1842)

Clinical Features of Bell's Palsy

Patients with Bell's palsy typically present with sudden onset (usually over hours) of unilateral facial paralysis, common findings include:

- Absent of forehead wrinkles “more by looking upwards”.
- Inability to close the eye.
- Absent nasolabial fold.
- Drop (asymmetrical) of mouth angle.
- Deviation the mouth to the non affected side.
Intact side

Very good note: that the patient may face a problem that he is can't taste in the affected side because of affection of Chorda tympani nerve that is the most distal branch intracanal so it is close to stylomastoid foramen and it is reversible damage.

Ramsay-Hunt Syndrome


Ramsay Hunt syndrome: acute peripheral facial neuropathy associated with erythematous vesicular rash of the skin of the ear canal.

Etiology: varicella-zoster reactivation.

Clinical picture:


- Vesicular rash of the ear or mouth.
- Ipsilateral peripheral facial palsy.
- Vertigo and ipsilateral hearing loss.
- Tinnitus.
- Otalgia: ear pain.
- Headaches.
- Fever.



A**B**

Inability to close the left eye
Mouth deviated to Rt non affected side (intact side)
No forehead wrinkles on the left side
No nasolabial fold in the Lt side

Bells palsy

C

- ① LMNL upper motor neuron lesion → C.B.T damage
- ② LMNL lower motor neuron lesion → nuclear or nerve lesion


- ① If damaged result contralateral lower part paralysis
- ② If damaged result ipsilateral hemifacial paralysis

In general the C.B.T gives dorsal and ventral contralateral in addition to dorsal ipsilateral

Dorsal part cause its importance for the eye has dual innervation from both cerebral hemispheres.

D → dorsal (anterior) part of nucleus for upper part of face

V → ventral (posterior) part of nucleus for lower part of face



upper motor neuron damaged at right cerebral hemisphere the defect will be contralateral lower (ventral) part
if the nucleus damaged or the nerve it will cause ipsilateral lower (ventral) part

في الحالة تبتع ال ٣ صور الي ب اخر سلايد، بما انو التأثير جاي على نص القيس اليسار معناها انو المشكلة بال LMNL وهاي
احنا عارفينه بتيجي ب defect اما بالنucleus or nerve السؤال الخلو انو كيف اميز بينهم ؟؟؟؟؟؟؟

Damaged Lt nucleus cause → Lt LMNL of facial nerve

Damaged Lt facial nerve → the same

How do we differ in between

We should remember cortico spinal tract the motor tract come from Lt and at lower part of MO cross to Rt

If stroke happened at Lt pons → ipsilateral facial palsy and contralateral hemiplegia

→ so Lt pontine lesion


If the nerve is damaged it will be without hemiplegia. Cuz the nerve out of brainstem

The Vestibulo-Cochlear Nerve (VIII) (Auditory or Stato-Acoustic nerve)


- It is a special sensory (hearing and balance) nerve.
- It emerges from the anterolateral surface between the pons and medulla oblongata with the facial nerve.
They are very closed together
- **Function:** Hearing and balance.
- **Modality:** Special somatic ^{Sensory} afferent.
- **Foramen:** Internal auditory meatus.
- It has two divisions:
 - Cochlear division.
 - Vestibular division.

Brainstem

Anterioinferior View



Note: Arrows indicate course of sound waves




The Glossopharyngeal Nerve (IX)

- The glossopharyngeal N. is a ¹⁰⁹⁷⁵mixed nerve containing:
 - a. Sensory fibers. Palat palatine tonsil pharynx carotid (body and sinus) post 1/3 of the tongue
 - b. Parasympathetic fibers. Give a branch for otic ganglion that supplies parotid gland
 - c. Motor fibers. Supplies one muscle: stylopharyngeous muscle
- It is more important as ^{mainly}sensory than as a motor nerve.
- **Origin:** from posterolateral sulcus of medulla.
- **Modalities:** GVE, SVE, GSA, & SVA.
- **Foramen:** Jugular foramen.✓

Brainstem

Anterioinferior View




Nuclei of the Glossopharyngeal Nerve

It has three nuclei:

9th
10th
11th

Ambiguous nucleus

1. The upper part of nucleus ^{مَشْرُوعِي 3 نِيرَق} ambiguous: it lies in the medulla (**motor function**).
2. The inferior salivary nucleus: it lies in the medulla (**parasympathetic function**).^{For parotid gland}
3. The solitary nucleus: it lies in the medulla (**sensory function**).^{Mentioned before}





Functions of the Glossopharyngeal Nerve

1. Sensory function: supplies the:

- a. Posterior 1/3 rd of the tongue (taste and general sensation).
- b. Mucous membrane of the pharynx and tonsil.
- c. Carotid sinus and carotid body.

2. Parasympathetic function: supplies the parotid gland.

3. Motor function: supplies one muscle only the stylopharyngeus muscle.



Lesion of the Glossopharyngeal Nerve

1. Loss of taste and general sensation from the posterior 1/3 of the tongue.
2. Loss of sensation in the pharynx If we touch its back it will cause gag reflex.
3. Ipsilateral loss of pharyngeal reflex Gag reflex
4. Some pharyngeal weakness (dysphagia).

Efferent (motor) vagus and cranial accessory

Afferent (sensory) glossopharyngeal

Evoked by touching the root of the mouth back of the tongue and prepare you for vomiting

Not complete cause it just supply one muscle

→ stylopharyngeous muscle

2 types of dysphasia:

Neurogenic and obstructive Called solid

1st → difficulty in fluid


2nd → difficulty in solid matters

The Vagus Nerve (X)

- The vagus nerve is the longest of the cranial nerve.
- **Origin:** from the posterolateral sulcus of the medulla.
Lower to 9th and lateral to 12th
- ❖ **It is a mixed nerve with:**
 - a. Motor fibers.
 - b. Parasympathetic fibers
 - c. Sensory fibers.
- ❖ It is the most important parasympathetic nerve in the
 - body.
 - **Modalities: GVE, SVE, GSA, SVA & GVA.**
 - **Foramen:** Jugular foramen.✓

Brainstem

Anterioinferior View



Nuclei of the Vagus Nerve

It has three nuclei:

- The main parasympathetic nucleus in the body


1. **Dorsal motor nucleus of vagus “Parasympathatic”:**

Gives parasympathetic fibers to respiratory system, digestive tract and the heart.

In GIT system all the system take from vagus until it reach the junction between proximal 2/3 of transverse colon and distal 1/3 and the distal to the junction is innervated by sacral
2. **Main part of nucleus ambiguous “Somatic Motor”:**

Gives motor fibers to the muscles of the larynx, pharynx and palate.
3. **Solitary nucleus “Sensory”:**

Receives taste and general sensations from the epiglottis and general sensation from the areas supplied by the parasympathetic fibers.



Function of the Vagus Nerve

1. The parasympathetic fibers supply:


- a. The smooth muscles and glands digestive system.
- b. The smooth muscles and the glands of the respiratory system.
- c. The heart.


2. The motor fibers supply:

- a. All the muscles of the larynx.
- b. All the muscles of the pharynx except the *Glossopharyngeal* **stylopharyngeus M.**
- c. All the muscles of the palate except the **tensor palati muscle.**
Otic ganglion by mandibular branch of 5th nerve

3. The sensory fibers:

- a. Receive sensation from the mucous membrane of the respiratory and digestive system mainly.
- b. Receives taste sensation from the epiglottis.
- c. Skin of the back of the auricle, external auditory meatus, lower and posterior part of the tympanic membrane.
- d. Meninges of the posterior cranial fossa.





Lesion of the Vagus Nerve

1. Palatal paralysis:

- a. Nasal tone of speech.
- b. Drop of palatal arch on paralyzed side and shift uvula to opposite side.
- c. Fluid regurge.
- d. Loss of palatal reflexes.

2. Pharyngeal paralysis:

- a. Neurogenic Dysphagia.
- b. Loss of pharyngeal reflex.

3. Laryngeal paralysis: hoarseness of the voice.

Others: tachycardia and constipation.


Sympathetic effect

The Accessory Nerve (XI)

- It is a purely motor nerve and has two roots:
 1. **The cranial root:** arises from the medulla oblongata. Joins the vagus nerve and distributed with its pharyngeal and laryngeal branches.
 2. **The spinal root:** arises by the upper 4-6 cervical segments of the spinal cord from the AHCs.
Anterior horn cells
- Supplies two muscles:
 - a. **Sternocleidomastoid muscle:** turn the head to opposite side.
 - b. **Trapezius muscle:** shoulder shrugging.
- **Origin:** from the posterolateral sulcus of the medulla.
- **Modality:** General somatic efferent.
- **Foramen:** Jugular foramen.
After they pass the foramen they divided another time to spinal and cranial the 1st supplies 2 muscles and the 2nd give vagus nerve.

Accessory Nerve

Schema



Cause the cranial mentioned
in motor branch of the vagus

Lesion of the Spinal Accessory Nerve

- Ipsilateral paralysis of sternocleidomastoid muscle (rotation of the head and neck to the same side). Head to site of lesion
- Ipsilateral paralysis of trapezius muscle (inability to shoulder shrugging).

The Hypoglossal Nerve (XII)


(The motor nerve of the tongue)

Supplies geniohyoid thyrohyoid and give branch to ansa cervicalis that connect it wit C2 + C3


- **Origin:** from the anterolateral sulcus of medulla oblongata. It receives a communicating branch from C1. *Motor*
- **Modality:** General somatic efferent.
- **Foramen:** Hypoglossal canal. *Specific foreman*
- **Function:** the hypoglossal nerve is a purely motor nerve which supplies:
 1. All the internal muscles of the tongue
 2. All the external muscles of the tongue: **hyoglossus, genioglossus and styloglossus, except the palatoglossus** (soft palate muscle supplied from the pharyngeal plexus). *vagus and cranial accessory*

Brainstem

Anterioinferior View



Hypoglossal Nerve Schema




The Spinal Roots and Nerves

- **Each segment has four roots:** two posterior & two anterior.
 1. **Two posterior (dorsal) roots:** there are sensory fibers (afferent fibers). Each posterior root has posterior spinal root ganglion.
 2. **Two anterior (ventral) roots:** there are motor and autonomic fibers (efferent fibers).
- The anterior and posterior roots will join to form a single mixed nerve called **the spinal nerve** → consist of 31 pairs of spinal nerves (8 cervical, 12 thoracic, 5 lumbar, 5 sacral, and 1 coccygeal).
- Each spinal nerve emerges from the intervertebral foramen.

The Exit of Spinal Nerve Roots

- ❖ Through the intervertebral foramina.
- ❖ Roots C₁- C₇ pass **above** their vertebra.
- ❖ C₈ pass **below** C₇ vertebra.
- ❖ Thoracic, lumbar and sacral nerves pass **below** corresponding vertebra.
- ❖ Lower lumbar, sacral and coccygeal nerves descend vertically as **cauda equina** pass below corresponding vertebra.

Relation of Spinal Nerve Roots to Vertebrae



Special Considerations of the Spinal nerve Roots

- **The first cervical nerve** and the **coccygeal nerve** usually have neither the posterior (sensory) roots nor the corresponding dermatomes.
- **The first cervical nerve** passes between the atlas and the skull.
- **The second cervical nerve** passes between the atlas and the axis.

Components and Branches of Spinal Nerves

- **The spinal nerve:** is formed by the union of posterior and anterior roots within the intervertebral foramen, resulting in a mixed nerve.
- **Posterior root.**
- **Spinal ganglion:** located within the posterior root and contains pseudounipolar neurons “First order neuron for somatic sensations”.
- **Anterior root.**
- **Spinal nerve rami.**

The Spinal Nerve Rami

- **Posterior ramus:** innervates the skin and muscles of the back.
- **Anterior ramus:** innervates the anterior and lateral muscles and skin of the trunk, extremities, and visceral organs.
- **Meningeal ramus:** innervates the meninges and vertebral column.
- **White communicating rami:** contain myelinated preganglionic autonomic fibers.
- **Gray communicating rami:** contain unmyelinated postganglionic autonomic fibers.

Functional Components of Spinal Nerve Fibers

1. Afferent Fibers:

A. General Somatic Afferent (GSA) Fibers: Convey **sensory input** from skin, muscle, bone, and joints to the central nervous system (CNS).


B. General Visceral Afferent (GVA) Fibers: Convey **sensory input** from visceral organs to the CNS.


2. Efferent Fibers:

A. General Somatic Efferent (GSE) Fibers: Convey **motor output** from anterior horn motor neuron to skeletal muscle.

B. General Visceral Efferent (GVE) Fibers:

- Convey **motor output** to glands, smooth muscle, and visceral and pelvic organs.





Visceral Nervous System

- 1. Visceral components of the spinal nerves:** are not present at all levels: they are associated mainly with T1-T2 and S2-S4 regions
- 2. Sympathetic Nervous System:** located within the lateral horns of T1-L2 segments of the spinal cord.
- 3. Sacral Parasympathetic Nervous System:** located within the anterior horns of S2-S4 segments of the spinal cord.

Spinal Nerve Innervation

- 1. Dermatome:** consists of **a cutaneous** area innervated by the fibers of **one spinal nerve**.
- 2. Myotome:** consists of **muscles** innervated by the fibers of **one spinal nerve**.
- 3. Sclerotome:** consists of **bones and ligaments** innervated by the fibers of **one spinal nerve**.

Thank You
THANK YOU

