


Lecture:

Autonomic Nervous System



Dr. Eyad M. Hussein
**Ph.D. of Neurology,
Consultant & Head of Neurology Department,
Nasser Hospital**

الرجاء تحويل الجوال إلى وضع الصامت
مع الشكر



Anatomy of the Autonomic Nervous System (Involuntary Nervous System or Visceral Nervous System)

- **Definition:** it is the involuntary nervous system which **supplies:**

- ✓ Unstriated (smooth) muscles of the viscera and blood vessels.
- ✓ Cardiac muscles.
- ✓ Glands.

- **Structures of the Autonomic Nervous System:**

- ✓ Sympathetic nervous system (Thoraco-lumbar)

"Fight or Flight".

- ✓ Parasympathetic nervous system (Cranio-sacral)

"Rest and Digest".

parasympathetic sympathetic

cranial part
Sympathetic
[Sacral part]

T₁ - L₂
or
C₈ - L₃

10973
[Brain stem]

S₁ - S₃

Parts of the Autonomic Nervous System

↳ has 2 order neuron.

in the CNS → first order neuron of ANS

Central part: consists of autonomic nuclei in the brain stem, thalamus, hypothalamus, insula and intermediolateral nuclei of the spinal cord.

↳ nuclei of the cranial nerves [10973]

in the PNS

2. **Peripheral part:** consists of: fibers and ganglia


• Peripheral Sympathetic nervous system (SNS).

• Peripheral Parasympathetic nervous system (PSNS).

* the first order neuron of the sympathetic → intermediolateral nuclei in the lateral horn cell [T₁-L₂]

→ fiber leave the 1st order neuron as preganglionic fiber to [symp. or parasymp. ganglia] 2nd order neuron
to reach the 2nd order neuron... Location of 1st order neuron

[white matter] Pre ganglionic + myelinated / ← 1st and 2nd order neurons (أبيض + غير ميالين) ←



Post ganglionic [gray matter] unmyelinated neur. • relay in the ganglia
↳ axon of 2nd order neuron.

The Origin of the Autonomic Nervous System

somatic-motor
for chest, abdominal
or resp. muscle.

Ant. horn cell T₁ - L₂
The origin of the sympathetic nervous system (Thoraco-lumbar nervous system): intermediolateral nuclei in the lateral horn cell ...

lat. horn cell between the
Ant. and post horn cell.

Consists of fibers which arise from the 12 thoracic and upper 2 lumbar segments (Thoraco-lumbar part) and reach the sympathetic chain by **the white rami**.


level S₂ to S₄ is
Ant. horn empty
PS nucleus

2. The Origin of the parasympathetic nervous system (Cranio-spinal nervous system):
the sacral part nuclei in Ant. horn cell of S₂-S₄ segment

a. The cranial part: consists of fibers which pass in four cranial nerves (3, 7, 9 & 10 cranial nerves)=**10973**.
nuclei of PS cranial N [10973]
the cranial nuclei → 1st order neuron of cranial part

b. The sacral part: consists of fibers which arise from **the S2, S3, and S4** segments of the spinal cord.

the PS nuclei in the Ant. horn cell [1st order neuron] of S₂-S₄



The Peripheral Autonomic Nervous System

↳ nerve or ganglia

- Most of the visceral structures are supplied by two systems (receive sympathetic and parasympathetic nerves).

in heart
↳ ↑ heart rate by sympathetic..
↳ ↓ heart rate by para sympathetic..

- The SNS supplies:**

1. Skin of the body (its arterioles, sweat glands).
2. Blood vessels of the limbs (its smooth muscles).
3. Deep viscera.

- The PSNS** ^{in the wall of the organs [target cell]} supplies the deep viscera only.

- Each of the two systems consists of two order neurons.

1st
2nd

مع الخوف مراح يوقف ① SNS
تقضي ال rectil villi muscle ②
pale face vaso const. by SNS [↓ BS in skin]

1. First Order Neurone

↳ inc NS → Brainstem
↳ Brain and spinal cord

- It is formed by:

1. The nerve cells of certain nuclei in the brain stem.
2. Lateral horn cells (intermediolateral nuclei) of D1-L2 and AHCs of S2-S4 of the spinal cord

- The axons of these cells carry stimuli to the autonomic ganglia (called preganglionic fibers which are white or myelinated).

axon
→ send the signal to 2nd order neuron.

* preganglionic fibers → long in parasympathetic
→ short in sympathetic

2. Second Order Neurone $\Rightarrow$ PNS

forming symp. chain $\leftarrow$ $\begin{matrix} \text{if} \\ \text{2} \end{matrix}$ $\begin{matrix} \text{in the ganglia} \\ \text{sup, inf} \end{matrix}$ $\begin{matrix} \text{Sympathetic} \\ \text{Parasympathetic} \end{matrix}$
 $\uparrow$ para vertebral or collateral ganglia $\leftarrow$

- It is formed by the nerve cells in **the sympathetic**

and parasympathetic ganglia. $\leftarrow$ terminal ganglia

- The axons of these cells carry stimuli to the

viscera (**called postganglionic fibers which are**

fibers begin after the ganglia

gray or non-myelinated). *myelinated* $\leftarrow$ عصب مغطى بغطاء

signal comes from general organ.

*يعني الاشارة تصل من عضو عام
heart beat lead to pain.*

*عضو خاص يرتبط به عصب
organ lead to pain.
 $\rightarrow$ sympathetic*

Types of Autonomic Ganglia

They are present outside the CNS and may be:

1. Paravertebral ganglia (Sympathetic): as sympathetic trunk (chain).
2. Collateral ganglia (Peripheral or Sympathetic): as celiac, superior mesenteric and inferior mesenteric ganglia.
3. Terminal ganglia (Parasympathetic): they are present on the walls of the viscera.




Notice that:

1. The sympathetic preganglionic fibers are **short**, while the postganglionic fibers are **long**.
2. The parasympathetic preganglionic fibers are **long**, while the postganglionic fibers are **short**.
3. The **preganglionic fibers** are white or **myelinated**.
4. The **postganglionic fibers** are gray or **non-myelinated**.

بالتوازن
Balanca

کی حفاظت ہے
Balance 2.

- Sympathetic Agonists
- Parasympathetic Antagonists



The body of the second neuron is found in the PNS and its axon is called **a postganglionic fiber**.

in the heart or GIT $\left\{ \begin{array}{l} \text{PSNS [stimulator]} \\ \text{SNS [inhibitor]} \end{array} \right.$

Transmitters of the Autonomic Nervous System

- ACh ← NT, ~50 Para Symp. = Symp. 10-20%
[nicotinic ACh]
1. The **preganglionic transmitter**:

For sympathetic and parasympathetic preganglionic transmitter is **Acetylcholine**.

2. The **postganglionic transmitter**:

A. All postganglionic fibers of the parasympathetic are

Acetylcholine.

muscarinic ACh
neuro muscular junction → ACh
but the somatic → nicotinic ACh.





B. The majority of sympathetic postganglionic fibers are


Epinephrine and/or Norepinephrine.

C. The few which are known to be **Acetylcholine** (for the sweat glands).

sweat gland 10-20%
Sympathetic
post gangl: 10-20%
Fiber is parasymp.
[ACh]

* occlusion in the PCA lead to Horner Syndrome → loss of sweat sensation
[ACh] parasymp. is to be NT, 10-20% Symp. ← [Anhidrosis]
pregang. postgang.





Key:


— = Preganglionic axons (sympathetic)
 - - - = Postganglionic axons (sympathetic)
 = Myelination
 — = Preganglionic axons (parasympathetic)
 - - - = Postganglionic axons (parasympathetic)

Comparison of Autonomic and Somatic Nervous System




Somatic <i>No relay.</i>	Autonomic <i>symp. parasymp.</i>
<u>Conscious</u> or <u>voluntary</u> regulation <i>منشئ مریکة، لا یرى</i>	Functions without <u>conscious awareness</u> (<u>involuntary</u>) <i>تصنیق بؤ بؤ یرى</i>
Fibers do not <u>synapse</u> after they <u>leave the CNS</u> (<u>single neuron</u> from CNS to effector organ). <i>[no ganglia]</i>	Fibers <u>synapse</u> once at a ganglion after they leave the CNS (<u>two-neuron chain</u>). Motor control.
Innervates <u>skeletal muscle</u> fibers, always <u>stimulatory</u>	Innervates <u>smooth muscle</u> , <u>cardiac muscle</u> , and <u>glands</u> ; either stimulates or inhibits

there is neuro muscular junction

between the muscle and the nerve.. [ACh nicotinic N.T] → may be blocked.



Comparison of Autonomic and Somatic Motor Systems

Cell bodies in central nervous system		Peripheral nervous system	Neurotransmitter at effector	Effector organs	Effect
SOMATIC NERVOUS SYSTEM		Single neuron from CNS to effector organs	 ACh	 Skeletal muscle	+ Stimulatory
		Heavily myelinated axon			
AUTONOMIC NERVOUS SYSTEM		Two-neuron chain from CNS to effector organs		 NE	  Smooth muscle (e.g., in gut), glands, cardiac muscle
		Lightly myelinated preganglionic axons	 Ganglion Unmyelinated postganglionic axon ACh Epinephrine and norepinephrine Adrenal medulla Blood vessel		
		Lightly myelinated preganglionic axon	 Ganglion Unmyelinated postganglionic axon ACh		

Organization of the Autonomic Nervous System

Characteristic	Sympathetic	Parasympathetic	Somatic*
Origin of <u>preganglionic</u> nerve	Nuclei of spinal cord segments <u>T1–T12</u> ; <u>L1–L3</u> (thoracolumbar)	Nuclei of cranial nerves <u>III, VII, IX, and X</u> ; spinal cord segments <u>S2–S4</u> (craniosacral)	
Length of <u>preganglionic</u> nerve axon	<u>Short</u>	<u>Long</u>	
Neurotransmitter in ganglion <u>pre</u>	<u>ACh</u>	<u>ACh</u>	
Receptor type in ganglion <u>pre</u>	<u>Nicotinic</u>	<u>Nicotinic</u>	
Length of <u>postganglionic</u> nerve axon	<u>Long</u>	<u>Short</u>	
Effector organs	<u>Smooth and cardiac muscle; glands</u> <i>same</i>	<u>Smooth and cardiac muscle; glands</u>	<u>Skeletal muscle</u>
Neurotransmitter in effector organs <u>post</u>	<u>Norepinephrine</u> (except <u>sweat glands</u> , which use ACh)	<u>ACh</u>	<u>ACh</u> (synapse is neuromuscular junction)
Receptor types in effector organs <u>post</u>	<u>α_1, α_2, β_1, and β_2</u>	<u>Muscarinic</u>	<u>Nicotinic</u>

Neurotransmission at adrenergic Neurons

● The process involves five steps:


1. Synthesis.
2. Storage. *uptake*
3. Release.
4. Receptor binding.
5. Removal of the neurotransmitter .
↳ and metabolism

بجده ساع يفرله
neurotransmitter

Pathway of Dopamine Metabolism

Synthesis

Degradation



1 SYNTHESIS OF NOREPINEPHRINE

- Hydroxylation of tyrosine is the rate-limiting step.

2 UPTAKE INTO STORAGE VESICLES

- Dopamine enters a vesicle and is converted to norepinephrine.
- Norepinephrine is protected from degradation in the vesicle.
- Transport into the vesicle is inhibited by *reserpine*.

3 RELEASE OF NEUROTRANSMITTER

- Influx of calcium causes fusion of the vesicle with the cell membrane in a process known as exocytosis.
- Release is blocked by *guanethidine* and *bretylum*.

5 REMOVAL OF NOREPINEPHRINE


- Released norepinephrine is rapidly taken into the neuron.
- Reuptake is inhibited by *cocaine* and *imipramine*.

4 BINDING TO RECEPTOR

- Postsynaptic receptor is activated by the binding of neurotransmitter.

6 METABOLISM

- Norepinephrine is methylated by COMT and oxidized by MAO.



Catecholamines A.A

- Catecholamines are sympathomimetic amines that contain the 3,4-dihydroxybenzene group such as:

1. Epinephrine. ✓

2. Norepinephrine. ✓

3. Isoproterenol. ✓

4. Dopamine. ✓

يعني خلاصة
epin. / norepin. ٢

Receptor Types in the ANS

1. Adrenergic receptors (adrenoreceptors):

↳ sympathetic

a. **Alpha Receptors:** are subdivided into two types:

- **Alpha “ α_1 ” Receptors:** are located on vascular smooth muscle of the skin, the gastrointestinal sphincters, bladder sphincters, and the radial “dilator” muscle of the iris.
 - ✓ **Effect:** produce excitation (e.g., contraction or constriction).
 - ↓ mydriasis, contraction of sph. → Bladder
 - constriction of BV. → GIT rectum
- **Alpha “ α_2 ” Receptors:** are located on walls of GIT, and wall of bladder.
 - ✓ **Effect:** often produce inhibition effect or relaxation.
 - wall of bladder, rectum ←
 - [relaxed in the rest]

Alpha 1 receptors



Vasoconstriction

↑BP

Increased peripheral
resistance

Increased blood
pressure



Mydriasis


dilatation of iris m.



Increased closure
of internal
sphincter of the
bladder

urination ↓, فزع
/ defecation.

Drugs Affecting Alpha Receptors



b. Beta Receptors: are subdivided into three types:

• **Beta “b₁” Receptors:** are located in the ^{heart} sinoatrial (SA) node, atrioventricular (AV) node, and ventricular muscle of the heart.

✓ **Effect:** produce excitation (e.g., increased heart rate, increased conduction velocity, increased contractility).

↑ cardiac output

• **Beta b₂ Receptors:** are located on ^{BV} vascular smooth muscle of skeletal muscle, ^{resp.} bronchial smooth muscle, and in the walls of the GIT and bladder.

↳ so may use in the asthma.

✓ **Effect:** produce relaxation (e.g., dilation of vascular smooth muscle, dilation of bronchioles, relaxation of the bladder wall).

• **Beta b₃ Receptors:** are located in the fat cells.

✓ **Effect:** Play important role with b₁ receptors in ^{help} lipolysis.

β -1 receptors

Heart

↑ heart rate

Increased
chronotropy and
inotropy

Increased
AV-node
conduction
velocity


Renal juxtaglomerular
cells

↑ renin production

Increased
renin release

↓
↑ angiotensin

β -2 receptors



Bronchial smooth muscle

resp.

→ spasm relaxation
Bronchodilation

we it use in bronchial asthma



Uterine muscle

Uterine relaxation
(tocolysis)

β -2 receptors




Bladder detrusor
muscle

Relaxation



Eye ciliar muscle

Relaxation



المعدة GI tract

Decreased
motility



Liver

Increased glucose
metabolism,
lipolysis



Smooth muscle

Relaxation

↳ parasympathetic.

a. **Nicotinic receptors:** are located in the **autonomic ganglia** (NN) of the **sympathetic and parasympathetic** nervous systems, at the **neuromuscular junction** (NM).

✓ **Effect:** produce usually **excitation** postsynaptic potential.

b. Muscarinic receptors: are located in the heart (M2), smooth muscle (M3), and glands (M3). *relaxation of rectum and bladder*

a. Are **inhibitory** in the **heart** (e.g., decreased heart rate, decreased conduction velocity in AV node). *↓ contractility function*

b. Are **excitatory** in **smooth muscle** and **glands** (e.g., increased GI motility, increased secretion).
 ↑salivation.

c. ^{direct} Are activated by ACh and muscarine.

d. Are blocked by atropine.

* the vagus nerve supply to the junction between R + 2/3 of transverse colon and L + 1/3 TC [Sacral N.]

muscle

a. A

d

b. A

G

c. A

d. A

Adrenal Medulla

← ganglion [symp.]
pre / long
post / short

- Is a specialized ganglion of the sympathetic nervous system.
- Preganglionic fibers synapse directly on **chromaffin cells** in the adrenal medulla.
- The chromaffin cells secrete **Epinephrine (80%)** and **norepinephrine (20%)** into the circulation.
- **Pheochromocytoma**: is a tumor of the adrenal medulla that
↳ malignant tumor.
secretes excessive amounts of catecholamines.
↳ ↑ HNT

AUTONOMIC

SOMATIC

Sympathetic innervation of adrenal medulla

Sympathetic

Parasympathetic

Preganglionic neuron

Ganglionic transmitter

Acetylcholine

Acetylcholine

Acetylcholine

No ganglia

Nicotinic receptor

Nicotinic receptor

Nicotinic receptor

Adrenal medulla

Postganglionic neurons

Neuroeffector transmitter

Epinephrine and norepinephrine released into the blood

Norepinephrine

Acetylcholine

Acetylcholine

Adrenergic receptor

Adrenergic receptor

Muscarinic receptor

Nicotinic receptor

Effector organs

Skeletal muscle

Autonomic Centers in the Brain Stem & Hypothalamus

1. Medulla oblongata:

- 
- Vasomotor center.
 - Respiratory center.
 - Swallowing, coughing, and vomiting centers.

3. Midbrain:

- Micturition center.

↳ anti-emetic act
in the vomiting center in
the medulla oblongata.

4. Hypothalamus:

- Temperature regulation center. + ↑ temp when we give him anti biotic [حارة يحمى] due to central fever.
- Thirst and hunger regulatory centers.

Effect of the Autonomic Nervous System on Organ Systems

Organ	Sympathetic Action	Sympathetic Receptor	Parasympathetic Action	Parasympathetic Receptor
Heart	$\uparrow$ heart rate $\uparrow$ contractility $\uparrow$ AV node conduction	β_1 β_1 β_1	$\downarrow$ heart rate $\downarrow$ contractility (atria) $\downarrow$ AV node conduction	M_2 M_2 M_2
Vascular smooth muscle	Constricts blood vessels in skin; splanchnic Dilates blood vessels in skeletal muscle	α_1 β_2	—	—
Gastrointestinal tract	$\downarrow$ motility Constricts sphincters	α_2, β_2 α_1	$\uparrow$ motility Relaxes sphincters	M_3 M_3
Bronchioles	Dilates bronchiolar smooth muscle	β_2	Constricts bronchiolar smooth muscle	M_3
Male sex organs	Ejaculation	α	Erection	M
Bladder	Relaxes bladder wall Constricts sphincter	β_2 α_1	Contracts bladder wall Relaxes sphincter	M_3 M_3
Sweat glands	$\uparrow$ sweating	M (sympathetic cholinergic)	—	—
Eye				
Radial muscle, iris	Dilates pupil (mydriasis)	α_1	—	—
Circular sphincter muscle, iris	—	—	Constricts pupil (miosis)	M
Ciliary muscle	Dilates (far vision)	β	Contracts (near vision)	M
Kidney	$\uparrow$ renin secretion	β_1	—	—
Fat cells	$\uparrow$ lipolysis	β_1	—	—

α_1	<p>Reflect the "flight or fight" RX.</p> <ul style="list-style-type: none"> Vasoconstriction of blood vessels (control of B.P.). Inhibit motility in the gut by contracting sphincter muscles and relaxing non – sphincter tissue. Mobilize energy by breaking down liver glycogen to glucose.
α_2	<ul style="list-style-type: none"> Found in presynaptic membranes. Provide feedback control of neurotransmitter secretion (inhibit Ca^{++} influx, decrease neurotransmitter release)(autoreceptor negative feedback).
β_1	<ul style="list-style-type: none"> Well known for their effects in the heart increase heart rate and force of contraction Induce muscle relaxation in the gut.
β_2	<ul style="list-style-type: none"> Induce bronchodilation. Induce smooth muscle relaxation in the gut. Induce conversion of glycogen to glucose. Stimulate secretion of insulin from pancreas.
M	<ul style="list-style-type: none"> Decrease heart activity. (M2) Increase motility in G.I. tract. (M1, M3) Depolarization of smooth muscle fibers, (M3) Hyperpolarization of cardiac muscle fibers. (M2)

Drugs Affecting Adrenergic and Cholinergic System

1. **Sympathomimetics Drugs:** *activate the Sympathetic* stimulation of adrenergic receptor.
2. **Sympatholytic Drugs:** *↳ ↓ NT* block the action of the neurotransmitters at the adrenergic receptors.
3. **Cholinergic “Anticholinesterase” Drugs:** stimulation the cholinergic receptors.
4. **Anticholinergic Drugs:** block cholinergic receptors.

The Parasympathetic Ganglia in the Head and Neck

There are four parasympathetic ganglia in the head and neck:

1. Ciliary ganglion. ✓
ocular motor nerve
2. Submandibular ganglion. ✓
3. Sphenopalatine (pterygopalatine) ganglion. ✓
4. Otic ganglion. ✓
glossopharyngeal nerve.
facial nerve via chorda tympani N.

*Sensory, motor Fiber just pass through ganglion and don't make relay in the ganglion..

Ciliary Ganglion

2nd order neuron of cranial nerve

Type: it is a small parasympathetic ganglion (1-2 mm in diameter).

↳ make synapse in ganglion

[Pre ganglionic → post ganglionic]

Site: in the posterior part of the orbit.

3 Roots Entering the Ciliary Ganglion: Just one make relay [para sy]

para and give post gang. fibers

1. Sensory root: branch from nasociliary nerve (branch of ophthalmic nerve). Just pass [not relay]

2. Sympathetic root: from the sympathetic plexus around the ICA. not relay / just pass

3. Parasympathetic root: from nerve to inferior oblique muscle (branch of oculomotor nerve). ↳ EWN.

make relay in the sup sympathetic ganglion [in the neck]

so through the ganglion pass post ganglionic fibers of sympathetic..

after the ganglion relay ↳ 1st order neuron give mixed fibers.. [sensory - symp. - parasymp.]


Branches of the Ciliary Ganglion

The ciliary ganglion has **8-10 short ciliary nerves** which pierce the sclera around optic nerve.


They contain:

1. **Sympathetic fibers:** to the blood vessels of the eye.
2. **Sensory fibers:** to the cornea and iris.
3. **Parasympathetic fibers:** to the supply the sphincter pupillae (for light reflex) and ciliary muscle (for accommodation).


بجسم
حواس
target
organ.



parasympathetic



Ciliary Ganglion Schema



* the short ciliary nerve carry post ganglionic parasympathetic fibers.

* the oculomotor nerve carry pre ganglionic parasymp. fiber.

* nerve of oblique M. carry pregang. parasymp. fibers

Submandibular Ganglion


- **Type:** it is a small parasympathetic ganglion (2-3 mm in diameter).
- **Site:** it lies in the submandibular region on the upper part of hyoglossus muscle.
- **Roots Entering the Submandibular Ganglion:**
 1. **Parasympathetic root:** *to the secretomotor to glands.* arising from the chorda tympani *preganglionic* nerve (Branch of facial nerve).
 2. **Sympathetic root:** *to the BV* arising from the sympathetic plexus around facial artery. *from plexus around ECA.*
 3. **Sensory root:** *to the sensory R.* from lingual nerve (branch of mandibular nerve).


Branches of the Submandibular Ganglion

1. Several small branches (**mixed: parasympathetic + sympathetic + sensory fibers**) pass directly from the ganglion to supply the submandibular salivary gland.
target organ
2. **Similar branches** join the lingual nerve to reach the sublingual salivary gland.
target organ.
of parasymp.

Mandibular Nerve [V3]


Lateral View





Sphenopalatine (Pterygopalatine) Ganglion

- **Type:** it is the largest parasympathetic ganglion (4-5 mm in diameter).
- **Site:** it lies in the pterygopalatine fossa.
- **Roots Entering the Sphenopalatine Ganglion:**
 1. **Sensory root:** two sensory branches arising from the maxillary nerve.
 2. **Sympathetic root:** the deep petrosal nerve from sympathetic plexus around the ICA.
 3. **Parasympathetic root:** the greater ^{superior} petrosal nerve (branch of facial nerve).
make relay.
GSPN



Branches of the Sphenopalatine Ganglion

Facial N → GSPN → the fibers relay in the Sphenopalatine + deep petrosal give post ganglionic < sympathetic > and bind with maxillary nerve to reach the zygomatic N → zygomatic temporal → communicate with lacrimal nerve of ophthalmic [CN V] → give < sympathetic > fiber to reach the lacrimal gland.


1. **Orbital branches:** enter the orbit through the inferior orbital fissure. They supply the orbital periosteum and the mucous membrane of the sphenoid air sinus. Another fibers pass with the zygomaticotemporal nerve to the orbit, where they branch off to join the lacrimal nerve to supplies **lacrimal gland**.

لacrimal gland (الغدة الدمعية) - من الأعصاب العينية.

Sphenopalatine division gland (الغدة الدمعية) sensory doesn't supply the gland.

2. **Pharyngeal branch:** to supply the mucous membrane of the nasopharynx.
3. **Greater palatine nerve:** to supply the mucous membrane of the hard palate and part of the nose.
4. **Lesser palatine nerve:** to supply the mucous membrane of the soft palate and the palatine tonsil.
5. **Short sphenopalatine nerve:** to supply the mucous membrane of the upper part of the lateral wall of the nose and nasal septum.
6. **Long sphenopalatine nerve:** to supply the mucous membrane of the anterior part of the hard palate.

Ophthalmic and Maxillary Nerves



Otic Ganglion

- **Type:** it is a small parasympathetic ganglion.
- **Site:** in the infratemporal fossa just below foramen ovale.
- **Roots Entering the Otic Ganglion:** *no sensory.*
 1. **Motor root:** arising from nerve to medial pterygoid muscle (*branch of the main trunk of mandibular nerve*).
 2. **Parasympathetic root:** formed by lesser superficial petrosal nerve (*tympanic / lesser petrosal n.* branch of glossopharyngeal nerve).
 3. **Sympathetic root:** from the plexus around middle meningeal artery (*from the plexus around ECA*).

* first order neuron $\rightarrow$ inferior salivary nucleus.

Branches of the Otic Ganglion

* the tympanic N of glossopharyngeal, lesser superficial petrosal N
is :- pre ganglionic fiber.

1. **Motor branch:** to tensor palati muscle and another branch to tensor tympani muscle.

2. **A communicating branch:** to auriculotemporal nerve,
this branch is formed of:


Handwritten notes:
- Sensation: $\rightarrow$ parotid gland (ATN) $\rightarrow$ parotid gland (ATN)
- Symp. $\rightarrow$ parasymp.
- Post. division of mandibular $\rightarrow$ post ganglionic fiber


a. **Sympathetic fibers:** to supply the blood vessels of parotid gland.

b. **Parasympathetic fibers:** secretomotor to parotid gland.

* the 2nd order neuron $\rightarrow$ otic ganglion


* the target organ :- parotid gland.





Mandibular Nerve [V3]

Medial View



The Parasympathetic Ganglia in the Head and Neck

Ganglion	Nucleus	Parasympathetic root	Sensory or motor root	Sympathetic root	Organs supplied
Ciliary	Edinger-Westphal	Oculomotor nerve int. oblique M.	Nasociliary V	Nasociliary nerve from internal carotid plexus ICA	Sphincter pupillae and ciliary muscles
Sphenopalatine	Superior salivary	Greater superficial petrosal nerve from VII nerve	Maxillary	Deep petrosal nerve from internal carotid plexus	Lacrimal glands; glands of nose, palate, mouth and pharynx
Submandibular	Superior salivary	Chorda tympani nerve from VII nerve	mandibular Lingual	Plexus around facial artery ECA	Submandibular and sublingual glands
Otic	Inferior salivary	Lesser superficial petrosal nerve from IX nerve	medial petrosal M. Mandibular "Motor"	Plexus around middle meningeal artery ECA	Parotid gland
Many ganglion in the different organs	Dorsal motor nucleus of vagus	Vagus nerve → many branches of vagus:- preganglionic → many para symp. ganglia:- 2nd order efferent in periph. organ.	Many nerves	Many plexus	Heart, Respiratory system, Abd. viscera or gland

Oculomotor

Edinger Westphal nucleus

EWN

3

oculomotor

inferior division of 3 nerve

nerve to inferior oblique M.

ciliary g.

s.c.n.

ciliary muscle

sphincter pupillae

Facial

superior salivary nucleus

SSN

7

facial

greater superficial petrosal n.

spheno-palatine g.

lacrimal gland

glands of nose & palate

chorda tympani n.

joins lingual n.

sub-mandib. g.

sub-lingual gland

sub-mandib. gland

Glossopharyngeal

inferior salivary nucleus

ISN

9

glosso-pharyngeal

tympanic branch

tympanic plexus in middle ear

lesser superficial petrosal n.

otic g.

auriculo-temporal n.

parotid gland

The Sympathetic Nervous System

- ❖ **Cells of Origin:** The sympathetic NS formed by the cells of the intermediolateral nuclei of the lateral horn of T1-L2 segments of the spinal cord (**Thoraco-lumbar part**) and reach the sympathetic chain by the white rami (right and left).
or maybe C8 - L3
- ❖ Sympathetic cells may also be found in C8 segments of the spinal cord.



The Sympathetic Ganglia

1. **Paravertebral Ganglia:** which formed the sympathetic chain or trunk.

18. 9, 10, 11, 12, 22-23 ganglia

- ❖ There are two sympathetic chains (right and left) on each side of the vertebral column.
- ❖ Each chain extends from the level of the **first cervical vertebra** (above) to front of **the coccyx** (below) where two chains unite to form a single coccygeal ganglion (**impar ganglion**).
- ❖ It formed of **22-23** ganglia on each side.

2. **Collateral or Peripheral Ganglia:** these are ganglia that surround some large arteries: **celiac, superior and inferior mesenteric ganglia**.

How Many Sympathetic Ganglia are Found in the Sympathetic Chains?

* there is 14 symp. segment but 22 paravertebral ganglion Two :-

① ← pre ganglionic fiber ← مقطع من relay بجمل white ramus ← ramus gray ramus (symp.)

1. **Cervical ganglia: 3 pairs.**

② ← lower fiber بعض lower fiber بعض symp. ganglia.

2. **Thoracic ganglia: 10-12 pairs.**

③ ← upper thoracic fibers هناك upper thoracic fibers بعض relay in sup., inf., middle

3. **Lumbar ganglia: 4-5 pairs.**

cervical symp. ganglia.

4. **Sacral ganglia: 4-5 pairs.**

④ ← fiber هناك fiber هناك relay in the collateral


1. **Coccygeal ganglion: one fused together**

18/19 ←

(impar ganglion).

to form single ganglion called impar ganglion

* the inf. cervical symp. ganglion fuse with the first thoracic and called stellate ganglion...





sympathetic ganglion


first rib


second rib


sympathetic chain

fourth rib









The Cervical Part of the Sympathetic Chain

Site: It lies behind the common, external, and internal carotid arteries, outside carotid sheath and medial to the vagus nerve.


It has:

1. **Preganglionic fibers:** from upper thoracic segments (T1-T4) until the three cervical ganglion. *↳ not relay. but ascend to cervical*
2. **Postganglionic fibers:** leave these ganglia, to the cervical nerves, cranial nerves and large vessels. *as EC A, IC A, subclavian A and it's branches.*

The Cervical Sympathetic Ganglion

There are three cervical sympathetic ganglion:

1. ^{larger} Superior cervical sympathetic ganglion.
2. Middle cervical sympathetic ganglion.
3. Inferior cervical sympathetic ganglion.
↳ fuse with first thoracic forming stellate ganglion.




Destruction of the Superior Cervical Sympathetic Ganglion

❖ This removes all the sympathetic fibers that reach face and neck and produces **Horner's syndrome**:

1. Myosis.
2. Partial ptosis.
3. Anhydrosis.
4. Enophthalmus.

not complete :- greater superioris
مؤثرات الأعصاب من الأعصاب الجذبية والكلى لم تصل جميعها
Ocular motor, N. lead to complete ptosis




myositis
partial ptosis

Sympathetic Supply of the Pupil

Pathway: Fibers start from hypothalamus, descends through the brain stem and cervical part of the spinal cord to the lateral horn cells of **C8 – T2 segment** → preganglionic fibers → sympathetic chain → relay in the superior cervical sympathetic ganglion. The postganglionic fibers pass with carotid plexus (around ICA) → long ciliary nerve → **dilator pupillae muscle**.

lead to mydriasis ↙

Para sympathetic



Innervation of the Urinary Bladder and Rectum

They receive 3 types of nerves:

1. **Somatic innervation (Pudendal nerve)**: supplies the striated ^[voluntary] ~~skeletal~~ muscles of the external sphincter of the urethra and rectum.



2. **Sympathetic nerves**: branches from the pelvic plexuses. They ^{→ contraction of the sphincter and relaxation of the bladder wall and rectum..} supply the blood vessels of the urinary bladder and rectum.

3. **Parasympathetic nerves**: branches from S2, S3 & S4 segments of the spinal cord → pass in S2, S3 & S4 → pelvic plexuses. **They supply:**

- The muscles of the walls of urinary bladder and rectum ^{contraction.} **(contraction).**
- Inhibitory fibers to the internal sphincter ^{relax} of the urinary bladder and rectum **(relaxation).**

urination and defecation ←
تجبريد
urination and defecation.


Effects of Autonomic Nervous System on Organs and Body

Organ 	Sympathetic action 	Parasympathetic action 
Eye: • Pupil • Ciliary muscle	- Dilates - Relaxes	- Constricts - Contracts
Glands: • Lacrimal & nasopharyngeal glands • Salivary glands	- Reduces secretion  - Vasoconstriction	- Increase secretion - Increase secretion
Heart	- Heart rate & muscle contraction increased	- Heart rate & muscle contraction decreased
Lung: • Bronchial muscle • Bronchial arteries	so asthma patient we give him - <u>Dilates bronchi</u>  Sympathetic mimitic - Constricts	asthma / give him parasympathetic - <u>Constricts</u> bronchi - Dilates
GIT: • Muscle in walls • Muscle in sphincters • Glands	relax sphincter - Decreases peristalsis - <u>Contracts</u> wall - Reduce secretion	contract. sphincter - Increases peristalsis - Relaxes wall - Increases secretion
Kidney	- Decreased urine secretion	- Increased urine secretion
Urinary bladder:  • Bladder wall • Sphincter	- Relaxes detrusor m. - Contracts sphincter	- Contracts detrusor m. - Relaxes sphincter

Organ	Sympathetic action	Parasympathetic action
• Gall bladder	- Relaxation	- Contraction
• Liver	- Increased conversion of glycogen to glucose	- Stimulate release of bile
• Pancreas	- Inhibition of insulin and glucagon secretion	- Stimulation of insulin and glucagon secretion
• Male sex organs —	- <u>Ejaculation</u>	- <u>Erection</u>
• <u>Pineal gland</u>	- Increase <u>melatonin</u> synthesis and secretion	-
• Skin:		
• <u>Sweat glands</u>	- <u>Increases secretion</u>	-
• <u>Pilomotor muscle</u> <p>العضلات الرفرفة [sympathetic]</p> <p>with stress → ↑ aggregation [hyper viscous syndrome] [symp.]</p>	- Contraction	-
• Platelet	Increase aggregation	-
• <u>lipolysis</u>	- <u>Increase</u>	-
• Renin secretion	- Increase — lead to ↑ BP	-
• Mast cells: histamine	- Decrease — (↓ dopamine source use it in electrical shock)	-

Parasympathetic


Sympathetic



Thank You



glossopharyngeal nerve:-



* in the PSNs
 → the preganglionic:- long..
 → the post ganglionic:- short..

[sympathetic]
 ... para sympathetic 11 ...

adrenal gland and the lateral ganglia

celiac or inf + superior mesenteric ganglion

[sympathetic] 11 ...
 in sympathetic chain close to the spinal cord
 [pregan:- short]
 [post gan:- long]

Sympathetic innervation of pupil

-: slide 72 & 73

→ the symp. give the signal from C8 - T1 spinal cord → pre ganglionic relay at the same side → ascend to sup. cervic symp. ganglion → post ganglionic fiber around the ICA → give long ciliary nerve supply the dilator ocular muscle lead to mydriasis..