

# Tumors

Cont.

ENAS ABU EWAIDA

**Oligodendroglioma** from oligodendrocyte  $\Rightarrow$  Schwann cell in PNS  $\rightarrow$  spongia

- **Best prognosis** of glial tumors.
- **5-15%** of gliomas.
- **4-5 decade**.
- **White matter** of cerebral hemispheres.

## Pathogenesis

- **IDH mutation** and whole arm chromosomal codeletion 1p and 19q.
- With progression to anaplastic form loss of **CDKN2A**.
- **No EGFR amplification** (in contrast to high grade astrocytoma).
- **TERT promoter mutation** (as in glioblastoma).

## Morphology

- Often with **cyst**, micro (sand-like) calcification. *fluid filled space. Cystic degeneration. forming space [cyst]*
- Microscopy: rounded, regular nuclei, relatively evenly-distributed chromatin (in contrast to astrocytoma), **white clear spaces around the nucleus** (*clear halo*), very **thin-walled blood vessels** (**chicken-wire pattern**). *↑ poleo morphism, ↑ mitotic activity. [grade 2  $\rightarrow$  grade 3]*
- IHC: IDH mutation might be detected. *cell border  $\rightarrow$  ..Qip, cell border  $\rightarrow$  sand like calcification..*

**Anaplastic oligodendroglioma:** more severe, has more mutations meaning it is more resistance to therapy and has a lower survival rate.

glioblastoma:  $\rightarrow$  grade 4  $\rightarrow$  upgrading oligo...  $\rightarrow$  glioblastoma

## Pilocytic astrocytoma

- Grade 1. Benign, well demarcated.. there is clefting between the normal and tumor cells.
- More discrete borders.
- Relatively benign behavior.
- Children and young adults. [rare in elderly]
- Cerebellum and around III ventricle.
- KIAA1549-BRAF gene fusion/duplication.

### Morphology

- Well-demarcated and cystic with mural nodule. <sup>solid area</sup>
- Limited brain invasion. there is space
- Loose microcystic and dense fibrillar areas. } morphology
- Microscopy: cells are elongated and has multiple thin processes (like hair).
- Rosenthal pattern, which is an elongated eosinophilic structure (characteristic in slowly growing tumors).

## Ependymoma

- Close to ependymal cells lined spaces (ventricles and central canal).  
↳ make collection in specific site in the ventricles of BS and called choroid plexus.
- 1-2 decades:

Near the IV ventricle. in children.

5-10% in the primary brain tumors in this group.

- In adults:  
↳ in adult

The spinal cord is the most common location.

Frequently in the setting of NF2.

NF: neurofibromatosis. 2 subtype  
↳ in the adult.   
NF1  
NF2 ⇒ most frequent in adult..

## Pathogenesis

NF2 gene mutated in spinal cord tumors (adults).

Distinct sets of driver mutations not shared with those of astrocytoma.

## Morphology

- Solid non-infiltrative. as pilocytic astrocytoma.
- Cerebrum, cerebellum and spinal. in adult
- Microscopy: oval cells, rosette formation (both types), there is a space between the cells and the blood vessels in which the processes of these cells extend.

True rosette formation: cells are arranged around a lumen.

Pseudo rosette formation: cells are arranged around blood vessels.

**2 variants:**

- **Myxopapillary ependymoma**: usually in filum terminale. *in the spinal cord..*
- **Subependymoma**.

**Filum terminale:** an extension of the pia mater that extends from the terminal end of the spinal cord to the tailbone.

## Choroid plexus tumors

- Rare.
- Benign or malignant.
- Choroid plexus papilloma: *most frequent*

Most common in children; lateral ventricles.

In adults **IV ventricle**.

Hydrocephalus.

CSF is increased in:

- **Overproduction** caused by a tumor in choroid plexus especially benign.
- **Obstruction**. of junction between the intra ventricular and subarachnoid
- **Reduced absorption** like in some types of arachnoidal inflammation leading to fibrosis of the granularity responsible for absorption. in the granularity [subarachnoid]

inflammation مع التهاب meningitis التهاب السحايا fibrinous inflam. الالتهاب الفيبروزي

## Neuronal and glioneuronal tumors

is same diff.  $\downarrow$

- 
- 

↳ diff. to produce neuron + glial cell.

- Gangliogliomas.  
↳ aggregation of cell body
- Dysembryoplastic neuroepithelial tumor.

# Medulloblastoma

embryo, which is derived from neural tube. • Embryo

not differentiated so called blast.

- Embryonal primitive neoplasm.
  - 20% of pediatric brain tumors.
  - Exclusively in the cerebellum.
  - Predominantly in children (less than 10 yo).
- \* 20% of pediatric tumor → CNS tumor  
 \* 20% of Brain tumors → medulloblastoma in pediatric

\* 20% of pediatric tumor  $\rightarrow$  CNS tumor

\* 20% of Brain tumors  $\rightarrow$  medulloblastoma in pediatric

## Embryonal primitive cells have:

- Large nucleus, evenly dispersed chromatin.
- Scarced cytoplasm, high nuclear/cytoplasmic ratio.
- High mitotic activity.
- Cells are similar to the origin.

\* proliferation pathways:- has role in the cell division, by the signals..  
WT pathway  $\rightarrow$  ABC

## Pathogenesis

**SHH** (sonic hedgehog) pathway: The vermis or lateral cerebellum. <sup>tumor in:-</sup>

WNT pathway: tumor inhib

- Around the anterior IV ventricle.
- Or middle cerebellar peduncle.

## Morphology

- Often <sup>relatively</sup> well circumscribed. rapidly proliferated..
- May involve the leptomeninges (the arachnoid and the pia mater) which may lead to CSF dissemination.
- Microscopy: monotonous cells, even chromatin. embryonic cell
- Mitosis is very high.
- Homer Wright rosettes.
- Desmoplastic nodular variant: Almost always SHH activated. \*anaplasia
- Large cell anaplastic variant. crumpled chromatin + granularity + pleomorphism
- Subarachnoid/SF dissemination.

more aggressive

**Karyorrhexis**: A stage of cellular necrosis in which the nucleus fragments and its chromatin is distributed irregularly throughout the cytoplasm.

## Meningiomas

- Tumors of adults, grade 1.
- Arise from the arachnoid <sup>origin</sup> (not a CNS tumor).
- Usually attached to the dura. tumor, part of dura
- Along any point of the external surface <sup>بالإضافة إلى طي الدورا</sup>
- Or within the ventricular system (from the choroid plexus). <sup>folding</sup>
- Previous radiotherapy is a risk factor.

## Pathogenesis

- Common lesion in NF2. with ependymoma
- 50-60% of sporadic cases show mutant NF2 gene.

## Morphology

- Gross:

En plaque growth: instead of forming a mass, we see thickening of the dura. <sup>طبقة</sup>

[layers]

→ sand like calcification as the oligodendroglioma.  
Firm to gritty (psammoma), like the calcification in oligodendroglioma but in a concentric pattern. → الجرسية

• Microscopy:  
↳ eosinophilic background, concentric nice structure called psammoma body.  
the cells has gated

3 types:

- Meningothelial.
- Fibroblastic.
- Transitional.
- They are similar to each other so it doesn't matter.

Concentric wrapping (whorls).

Psammomatous: psammoma body, كتلة كبيرة من الجرسية

\*traditional meningioma → grade I [Benign] need surgical removal. ديسر الوفاة

### Atypical meningioma

- Grade 2.
- One fourth of the cases. 1/4 cases.
- Higher recurrence rate.
- RT in addition to surgery.
- Aggressive features.
- Clear cell and choroid patterns are grade II by definition.

### Anaplastic meningioma

- Grade 3.
- 1-3%. rare [more aggressive]
- Carcinomatous/sarcomatous features instead of the typical features of meningiomas.
- Papillary and rhabdoid variants.  
↳ grade 3

ببطل تبه د  
meningioma...  
ugly + sat