


CNS PATHOLOGY

Lecture I



CEREBROVASCULAR DISEASE

Cerebrovascular disease

- Ischemic or hemorrhagic etiologies

- Ultimately infarction → Tissue necrosis

- Stroke versus TIE → Transient ischemic attack

Cerebrovascular disease of CNS if last < 24 h

Cerebrovascular disease of CNS if last > 24 h

- The 3rd leading cause of death → After cardiovascular diseases and cancer

- Ischemia/hypoxia

Deficiency of blood supply

Deficiency of O₂ only

- Global or focal
- ✓ Manifestation related to the region affected
- ✓ Embolism more common than thrombosis

Global: if the entire CNS is affected
focal: only portions

This will result in neurologic findings

Ischemia

→ Deficiency of blood supply

→ O₂ deficiency +
nutrients deficiency +
deficient waste products removal


→ It's more hazard than hypoxia


Cerebrovascular disease

- Hemorrhage ^{خاصة ما يكون} **malignant HTN** → $>200/120 \text{ mmHg}$
 - Due to hypertension and vascular anomalies (aneurysm and malformation)

ربما لا تقاوم الجدار بضغط عالٍ
T. intima & T. adventitia
Cannot stand high BP

↓
Brain aneurysms &
Congenital aneurysms
← characterized by loss of
Tunica media






Hypoxia and ischemia

- 20% of O₂ consumption
- Autoregulation of blood flow
- Strictly dependent on aerobic metabolism

→ Although the CNS constitutes only 2% of body mass!

So it is very susceptible to ischemia



Hypoxia and ischemia

- The size of the lesion determined by

- Duration

- Magnitude and rapidity of the reduction

- Collaterals

*If gradually developed
(through months, years...)
there's a chance for
collateral circulations*

*Small BV will induce less
lesions comparable to the
lesions that will result
from large BV*

- From C. of Willis/external carotid/ophthalmic
- Cortical/leptomeningeal
- Deep parts lack collaterals

→ So more susceptible to infarctions

Hypoxia and ischemia

- Excitotoxicity

- Release of glutamate NT → very toxic ↓
- Damage neurons by excessive influx of Ca^{++}

طبیعی آبی neuron لا یعرضن لـ hypoxia و ischemia
بعض Release of glutamate

- Penumbra → Partial ischemia

- Area of "at risk" brain tissue
- Can be rescued by antiapoptotic interventions

Ischemia is potent inducer of cell death 'Apoptosis'
ربالتیج له استهتنا apoptotic mechanisms
هتقدر نتقنال neuron death

لا یحصل occlusion لـ artery هین
صیحصل infarction فی المنطقة
ابی تیغذ هاب فی مناطق متکون
جوانین های المنطقة بیهیه فی
نقصان لـ blood بس من
بس کل کل های هیه ار
Penumbra

Focal cerebral ischemia

■ Embolism

■ Sources

- Most commonly cardiac thrombi
- Atheroma plaque within the carotid
- Paradoxical
- Cardiac surgery
- Tumor, Fat, and air

هو من أكثر الأماكن affected
بـ Atherosclerosis وخاصة
عند bifurcation into
ICA & ECA

After trauma

- Most frequently the middle cerebral artery

تستمر من
of the Common
Carotid artery

In general, the most common cause of embolisms are DVTs

وفي DVT الهدف عادة يكون الجبهة اليمنى (RV)
يعني embolism into the lung
لكن انه يرجع على الجبهة اليسرى (LV)
ومن ثم إلى CNS أود أي مكانه تالفة عادة
بأصبع إلا إذا في defect في Heart
(Ceg. interventricular septum defect)

Focal cerebral ischemia

- Thrombotic occlusion

- Most commonly acute plaque changes in the bifurcation


- Inflammatory process

- Syphilis and TB
- Vasculitis

very rare

Disruption of the plaque → Rupture → Ulceration

↳ Bcs of slowing of BF (Turbulence)



central fat 'lipid-laden macrophages'

C.T. wall 'endothelial cells'


↳ If disrupted → exit of collagens
'become exposed to blood'

↓
Thrombus formation ← Collagen is strong inducer of coagulation

Focal cerebral ischemia

- Other

- Hypercoagulability
- Dissecting aneurysm
- Drug abuse




Focal cerebral ischemia

- Starts as nonhemorrhagic (pale/anemic)
- Secondary hemorrhage
 - Due to reperfusion → *Reperfusion injury* →
 - Leading to hemorrhagic infarct
 - Petechial morphology
 - Cautions with thrombolytic treatment ✓✓

لما منطقة تكون ischemic أو زرع
عليها الـ blood مستعمل
كمية الـ O₂ وبالتالي هتكون
free radicals من كيميائية


Secondary hemorrhage
due to ischemia
and reperfusion injury
هتكون على شكل نقاط لكن
لما آتت injury من ان وصل تكون
hemorrhagic هتكون بي
Thematoma



→ The infarcted area

→ Secondary petechial hemorrhage

B



Focal cerebral ischemia

- Gross morphology

- First 6 hours

- Little change

- By 48 hours

- Pale, soft swollen, indistinct gray-white matter

→ Discrimination btw white and gray matter will be lost

- 2-10 days


- Gelatinous, friable, distinct normal/infarcted area

→ The infarcted area become well demarcated from the surrounding healthy area

- 10 days-3 week

- Liquefaction, fluid filled cavity

↓
Type of necrosis



Focal cerebral ischemia

- Microscopic

- Acute infarct

- 6-12 hours

- Eosinophilic neuronal necrosis

- Edema

- Indistinct gray/white matter

- Disintegration of the glial cells, fibers and endothelium


- 48 hours

- Neutrophilic emigration ✓

Due to disintegration of genetic materials (DNA & RNA) which are stained with blue (Basophilic)

↳ So when disintegrated, the acidophilic components will constitute the largest portion of the cells

Bright eosinophilic pyramidal neurons → i.e. necrotic



Focal cerebral ischemia

- Microscopic

- Subacute (evolving) infarct

- 48-72 hours

- Phagocytes and microglia
 - Reactive astrocytes
 - New blood vessels (as early as 1 week after the insult)
 - Astrocytes at the edge develop a network of cytoplasmic extensions
- Then ()
- Then ()
- deep areas Repair outer areas
Start from outside toward the center
- Granulation tissue formation

Almost completely occupied by inflammatory cells
(mainly macrophages)

neutrophils & eos.
leukocytes
...

→ new blood vessels

B

Focal cerebral ischemia

■ Microscopic

■ Healed infarct (after several months)

- Dense meshwork of glial fibers → This will completely replace the infarcted area
- New capillaries
- Perivascular connective tissue
- Chronologic morphologic changes

نوع من 'Zoonation'
يعني المنطقة الأقرب لل
Periphery تكون
قطعت شوطاً كبيراً
في عملية Repair

■ Hemorrhagic infarction

نفس الخطوات الـ Ischemic
تُضاف إليها blood extravasation

- The same with blood extravasation added
- Extensive if under anticoagulant
- Venous infarction
 - Thrombosis of the sinuses

كل الـ Arterial infarction
يمكن تحويله لـ Venous

Cavity surrounded by
meshwork of glial
cells, BVs & CT
(gliosis)

Infarction
↖




Lacunar infarcts

Small infarcts due to occlusion of small BVs

- Occlusion of small vessels as a result of arteriolosclerosis/hypertension
- Less than 15 mm
- Single or multiple
- Involves, putamen, globus pallidus, thalamus. Internal capsule, deep white matter, caudate nucleus and pons
- Tissue loss surrounded by gliosis ✓

المنطقة
بنيّة
inner areas
(Basal ganglia)



Global cerebral hypoxia/ischemia

- Cardiac arrest, shock, and severe hypotension
- CO poisoning
- Transient postischemic confusional state with complete recovery
بعض ما نسميها oriented لكن بعد ما بغير الموضوع يرجع طبيعي
- Or irreversible damage; "diffuse hypoxic/ischemic encephalopathy"

Global cerebral hypoxia/ischemia

- Border zone (watershed) infarcts

- Border between arterial territories → مناطق بين البينيين، يتأثر من هنا
شعيرة Blood ومن هناك شوية
- Result in wedge shaped infarct
- With secondary hemorrhage
- Often bilateral
- Commonly after cardiac arrest resuscitation

نندجف اناء ال hemorrhage عى سكل
نقاط (Petechial hemorrhage)
مبالتالى ناجم من Ischemic
طيس hemorrhagic infarct




Global cerebral hypoxia/ischemia

- Morphology

- Edematous and swollen brain
- Poor gray white demarcation
- Irreversible injury

- 6-12 hours; dead red neuron
- 24 h-2 weeks, Subacute changes; necrosis, macrophages, vascular proliferation and gliosis
- After 2 weeks; removal of necrotic tissue and gliosis

← تَقْرِيبًا فُوكَال



Intracranial hemorrhage

- Epidural and subdural; mostly trauma related ①
- Subarachnoid and intraparenchymal; related to cerebrovascular disease ②


Intraparenchymal hemorrhage

- Sudden onset of symptoms
- Versus secondary hemorrhage
- Middle/late adult life

From Basal ganglia → Ganglionic versus lobar hemorrhage → from lobe 'outer surface (gray matter)' of the cortex


- Hypertension is the most commonly associated factor with ganglionic intraparenchymal hemorrhage
 - 50-60% Putamen

Proliferated arteriolosclerosis → malignant HTN
Hyaline arteriolosclerosis → HTN within the normal range



Intraparenchymal hemorrhage

- Effects of HT on blood vessels
- ☞ ■ Cerebral amyloid angiopathy is the risk factor most commonly associated with lobar hemorrhage
- Morphology → Hematoma and blood collection



landmarks of layers of
the vessel wall cannot
be identified
'arteriosclerosis'

This is a histological micrograph showing a cross-section of a small blood vessel. The vessel lumen is the clear central area. The vessel wall is thickened and stained a uniform pink color, which is characteristic of hyaline arteriosclerosis. The normal layered structure of the vessel wall (intima, media, and adventitia) is obscured by this uniform staining. Numerous dark blue, oval-shaped nuclei of surrounding cells are visible in the tissue.




So the type of arteriosclerosis
is hyaline arteriosclerosis

A

Subarachnoidal hemorrhage

- The most common cause is rupture of saccular "berry" aneurysm in a cerebral artery


خاصة في
C. of willis
where the
arteries are
branching


- Saccular aneurysm (نزى الكرش) 
- 2% of the population → عدد كبير نوعاً ما
- 90% found near major arterial branch points in the anterior circulation

Saccular

الناس الي يموتوا فجأة يكون
عندهم ضغط دم مرتفع جداً
يكون داء المنعرج aneurysm

The anterior circulation is the most frequently affected





Subarachnoidal hemorrhage

- Pathogenesis
 - A developmental anomaly
 - Absence of smooth muscle and elastic lamina
 - The majority are sporadic
 - Cigarette and HT
 - Genetic factors

Subarachnoidal hemorrhage

■ Morphology

Saccular ← *mms to cms*

- Shape and size

- Atheroma, thrombus → *Due to turbulence and slowing of the blood in the dilated part*

ما قبل الموت
autopsy
signs of prior
hemorrhage


- Prior hemorrhage

- Bleeding into the subarachnoid or brain substance

- Absence of smooth muscle and elastic lamina

- Thickened hyalinized intima and a covering of adventitia ✓

↓
i.e. T. media



Vascular malformations

- 4 groups

- AV malformations

→ Abnormal connection btw the artery and vein

- Risk of hemorrhage

- KRAS mutation *→ Maybe a type of neoplasm??*

- Cavernous malformations

extremely dilated BVs

- Risk of hemorrhage

- Capillary telangiectasias

- Venous angiomas

Vascular malformations

- Morphology

- AV malformations

→ Serious problem as it is a significant cause of intracranial hemorrhage

- Warm like

- Pulsatile shunting

artery ← لينة أمه
vein ← باض جوا
Pulsation ← فوكون في

- Subarachnoid or brain


- Prior hemorrhage → Dark brown due to hemosiderin accumulation


- Duplication and fragmentation of elastic lamina

- Thickening and replacement of media by hyalinized CT

- Separated by gliosis

↓
hemorrhage





Vascular malformations

- Morphology
 - Cavernous malformation
 - No brain parenchyma
 - Most often cerebellum, and pons
 - Low flow → non-pulsatile
 - Old hemorrhage

الخرف

Vascular dementia

→ dementia
loss of brain material
يستجى للناس الى أعمالهم تبتة

- Multiple infarcts over years
- Dementia, gait abnormality → Disorientation . . .
سنة غير طبيعية loss of motor function . . .
- Caused by
 - Atherosclerosis
 - Thrombosis/embolus from the carotid or heart
 - ✓ ■ Arteriolosclerosis from chronic HT