

Blood supply of the Brain

*Arteries of the Brain — 2 carotid [Ant.]
— 2 vertebral (post)

the Branches anastomose on the inf. surface of the base of the brain to form the **circle of willis**

Internal carotid Artery:-


origin :- at the bifurcation of the CCA

* ascend to the carotid canal of the temporal bone → cavernous sinus → enter the dura and arachnoid [medial to the anterior clinoid]

* Bifurcate to MCA and ACA at the level of end of the Sylvian fissure ...

ICA segments; -

- 1- cervical segment [C1]
- 2- petrous segment [C2] horizontal in the temporal bone
- 3- cavernous [C3]
- 4- intracranial (supra clinoid) [C4]. **has 3 parts :-**
ophthalmic - communicating - choroidal



ACA

MCA

* Smaller

* anterior-medial

* A com A. join the 2 ACA

* Branches; - **more**

anterior perforated sub. part of the lentiform and cuneate nuclei and the internal capsule

* larger (ICA continuation)

* laterally in the lateral cerebral sulcus [sylvian fissures]

* supply the lat. surface

* Branches; - **less**

anterior perforated sub. part of the lentiform and cuneate nuclei and the internal capsule

vertebral Artery:-



- branch of the 1st part of the subclavian artery

- at the upper 6 vertebrae pass through foramina

in the transverse process. [طابقه فم 7 C 7]

- enter the skull through the foramen magnum and pierce the dura mater and arachnoid

- pass upward, forward, medially around the medulla



ICA angiogram AP & Lat

- At the lower border of the pon, it joins of the opposite side to form the basilar artery.

VA branches:- [intracranial]

-meningeal branches :- small vessels.

-post. Spinal artery 2..

-Ant. spinal artery 1...

- PICA :- from Vertebral and Basilar and supply choro plexus of 4th V.

(injury in the PICA lead to wallenberg syndrome.. [vertigo-diplopia dysarthria, ipsilateral Horner syndrome, numbness, ipsilateral face and contralateral limb. , no limb weakness, no Loc]

- medullary branches.

Basilar Artery:- formed by union of 2 vertebral A.

- Ascend in a groove on the Ant surface of the pons.

- divided into 2PCA at the upper border of the pons.

Branches of MA.

1-Pontine A.

2- Labyrinthine A.

3- AICA

4- SCA.



5- PCA supply the visual cortex, post choroid plexus in inf. horn and 3rd v. and injury in this Artery lead to contralateral

hemianopia

circle of willis i10A: Acom.A, 2ACA, 2ICA, 2PcomA ,2PCA, basillar function:- Blood distribution and variation.

Function & Blood supply of Internal Capsule

في الـ 10A
in the Basal ganglia level



Supply	Area
Mainly by the medial and lateral striate central branches of MCA. ACA	Corpus striatum & internal capsule
P Com A, Basilar, PCA	Thalamus
PCA, Basilar, SCA	Midbrain
Basilar, SCA, AICA	Pons
VA, PICA, Basilar, Anterior & Posterior spinal art.	Medulla oblongata
SCA, AICA, PICA	Cerebellum

Basilar is formed by multiple Arteries and is a very important artery.

control of cerebral Bf.


$\uparrow \text{CO}_2$, $\uparrow \text{H}^+$, $\downarrow \text{O}_2$, parasympathetic $\rightarrow$ vasodilation

$\downarrow \text{CO}_2$, $\downarrow \text{H}^+$, $\uparrow \text{O}_2$, sympathetic $\rightarrow$ Vasoconstriction.

veins of the brain: has thin wall, no muscular tissue, no valves.

drain into the cranial venous sinuses.

Cerebral Autoregulation



spinal cord arteries: longitudinal..

- 2 post. Spinal A. (1/3): small in upper thoracic

- 1 Ant. spinal A. (2/3): - small in upper and lower thoracic.

* reinforced by small segmentally arranged A:- **great anterior medullary A. of Adamkiewicz** .. [لحمية منية ظهر راجع حبل الحبل]

↳ from the aorta, in the lower thoracic [T10-L2], unilateral, left side and supply the lower 2/3 of cord..

clinical Notes; -

- cerebrovascular disease: **50%** Of all adult neurologic hospital admission
- if the Brain Blood flow completely cut due to cardiac arrest for example lead to:-
 - 5-10 sec;-loss of conscious
 - 1min:- neuronal function cease
 - 4min: irreversible damage

Arterial occlusion

ACA occlusion	MCA occlusion	PCA occlusion
* sensory loss	* contralateral hemiparesis	* contralateral homonymous
* Contralateral leg and foot weakness	* hemisensory loss in face and arm.	hemianopia with macular sparing [occipital]
* personality changes	* Aphasia mostly if Lf. side affected	* impairment of memory [temporal]
[distal to A.com. A]	* contralateral homonymous hemianopia [optic radiation]	

ICA Occlusion	vertebral Artery occlusion
<ul style="list-style-type: none"> * contralateral hemiparesis * hemisensory loss in face and arm. * Aphasia mostly if Lf. side affected * contralateral homonymous hemianopia [optic radiation] 	<ul style="list-style-type: none"> * pain, temp. sensory loss [ipsilateral face, contralateral Body] * Hemianopia or complete cortical Blindness * ipsilateral loss of the gag reflex, dysphagia and hoarseness. * vertigo, nystagmus, nausea, vomiting, coma * ipsilateral Horner syndrome * unilateral or bilateral hemiparesis * ipsilateral ataxia and other cerebellar signs

cerebral Blood flow impairment

causes; → * disease that lead to alteration in BP [postural hypotension, heart disease....etc]

* disease of arterial walls [atheroma]

* disease due to Blockage of the arterial lumen [thrombus, fat globules]..

cerebral Aneurysms:- * dilatation of the arterial wall and may be ruptured .
* usually occur at branching points of major Arteries.

presentation :- pressure symptoms, subarachnoid hemorrhage

cerebral hemorrhage:- * rupture of the thin walled lenticulostriate arteries
* (most commonly) basal ganglia hemorrhage
* Damage to corticonuclear and corticospinal fiber,
* coma, contralateral hemiplegia, HTN