

Ventricular System and Cerebrospinal Fluid

Dr. Nidal Abuahdrous

Consultant Neurosurgeon

Certificate of Arab Board of Neurosurgery

Head of Neurosurgery at European Gaza Hospital

Ventricular System and Cerebrospinal Fluid

Objectives

- ^{in the ventricles} CSF location, function, production & absorption
- Blood-brain barrier
- Blood-cerebrospinal fluid barrier

Ventricular System

- 2 lateral ventricles

- Interventricular foramina (of Monro)

Join with 3rd ventricle

- 3rd ventricle

- Cerebral aqueduct (aqueduct of Sylvius)

in midbrain

- 4th ventricle

- Central canal of the spinal cord
- 3 foramina in its roof, with the subarachnoid space (2 Luschka, 1 Magendie)


↳ 4th ventricle

Post fossa outlet


Pons upper medulla

(brain stem)


*Pons ← 4th ventricle → cerebellum
medulla oblongata*



A



Connection
between 2 thulmi




Lateral ventricles → has 3 horn

- Body
- Anterior (frontal) horn
- Posterior (occipital) horn
- Inferior (temporal) horn
- Trigone (Atrium)

→ connect Body with occipital, temporal
 → common site of meningioma
 → choroid plexus papilloma
 ما بتيجي على ventricles
 كتير بتيجي على cortex
 أكثر (لكن لو مارضنا بكونه مش ممكن trigone رينا)

شنت
 شنت
 شنت
 hydrocephalus



3d & 4th ventricles

- 3d ventricle: a slitlike cleft between the two thalami
فتحة رقيقة
➤ Choroid plexus is suspended from the roof
معلقة به
CSF production.
- 4th ventricle: a tent-shaped cavity
➤ Situated anterior to the cerebellum and posterior to the pons and the superior half of the medulla
➤ Floor: a rhomboid-shaped نرى المعبر
➤ Roof: Superior medullary velum & Inferior medullary velum
➤ Choroid plexus is suspended from the inferior half of the roof (PICA)

Choroid plexus of the ventricular system

فوقه frontal horn و temporal horn و تحتيه body و تحتيه inferior horn
Monro و تحتيه roof of 3rd ventricle و تحتيه medial part و تحتيه

- In lateral ventricles:


- Body: medial & inferior
- Inferior horn: medial & superior

- In 3rd ventricle:

- roof


- In 4th ventricle:

- the inferior half of the roof
lower




Rt lateral Ventricle

(via transcallosal approach)



Transchoroidal approach to 3d Ventricle

- Transchoroidal approach directed through the body portion of the choroidal fissure using an opening through the corpus callosum
- The choroidal fissure has been opened by incising along the tenia fornicis, not tenia thalami



Subarachnoid space

- Between the arachnoid & pia mater, around the brain and spinal cord
- Filled with CSF
- Contains the large blood vessels of the brain
- Basal subarachnoid cisterns
- In Spine: it ends at S2-S3

in skull base

Subarachnoid space

منطقة محتوية لـ CSF

متفخة

→ the spinal cord ends in L1 - L2

Cerebrospinal Fluid

السائل النخاعي

Table 16-1

Cerebrospinal Fluid Physical Characteristics and Composition

Appearance	Clear and colorless
Volume Brain & spine	c. 150 mL
Rate of production	0.5 mL/min
Pressure (spinal tap with patient in lateral recumbent position)	60–150 mm of water
Composition	
Protein	15–45 mg/100 mL
Glucose	50–85 mg/100 mL
Chloride	720–750 mg/100 mL
Number of cells	0–3 lymphocytes/mm ³

[CSF] clear color less normally.

→ if there is obstruction in CSF pathways lead to cell death → ↑ ICP
 ← خلل في السائل ←

[hydrocephalus] in adult may lead

to urgent surgery—

adult die on one day
 child → surgery → يفتح السائل
 منقوع

زيادة infection جفاف infection

☆

عليا 200

in bacteria → neutrophils / viral → ↑

lymphocyte

CSF function

Table 16-2 Cerebrospinal Fluid Functions

1. Cushions and protects the central nervous system from trauma
2. Provides mechanical buoyancy and support for the brain
3. Serves as a reservoir and assists in the regulation of the contents of the skull
4. Nourishes the central nervous system
5. Removes metabolites from the central nervous system
6. Serves as a pathway for pineal secretions to reach the pituitary gland


atmosphere جاي
Brain ٻي
[mechanical protection]

regulation of the contents
provide some nutrition
atmosphere جاي

Hormones.


كل علكه الة كصاب من Pressure ↑ لا نه الرماخ لا يتحل الصفه العالي و بجوترا الانسان منه
 سواد كذا رسم اذ غير حانه حافظ على pressure intercranial

Monro-Kellie Doctrine



طه المصنفه الة كصاب من خلايا في neuro ← تضيق ال Press. de compression.

Volume-Pressure Curve



CSF formation

inside the ventricular system

main producer of CSF

- Mainly in the choroid plexuses


→ active secretion need transport and energy.

- Active secretion
- Not pressure regulated

active secretion

- Ependymal cells


- Brain



* production from blood to CSF

lateral V. $\rightarrow$ 3rd V. $\rightarrow$ 4th V. $\rightarrow$ Sub arachnoid space in brain (absorption is occure)

CSF circulation



Blood-Brain & Blood-CSF Barriers

← تمنع دخول المواد
الضرورية
إلى الدماغ

- The CNS requires a very stable environment In order to function normally
بحاجة تزيير لا تنقص

- This is provided by isolating the nervous system from the blood by these barriers

- They protect brain and spinal cord from potentially harmful substances while permitting gases and nutriment to enter the nervous tissue

blood + brain { blood + CSF


1 Blood-Brain Barrier (BBB)

Structure

- Endothelial cells
➤ Tight junctions (Lipid)
- Basement membrane
- The foot processes of the astrocytes

In newborn: BBB is more permeable (Bilirubin → kernicterus)

خلو صبار و عدم Junction
بيكون نفاذ فنتجنا اليرقان
في عكس الكبار Bilirubin لا ينفذ
BBB



Blood-Brain Barrier (BBB)

حالی ماذا تلتحق نفاذ به BBB

- BBB permeability is inversely related to the size of the molecules and directly related to their lipid solubility

ملاحظة معينة

کی ما کنت Solubility عدت بسهولة

کی ما کبر حجم الجزيء
قل نفاذيته عبر BBB

- BBB is absent in:


هنا
metastatic tumors

- pineal gland هنا يوصل hormones
- posterior lobe of the pituitary
- tuber cinereum
- the wall of the optic recess
- area postrema (sample the chemical content of the plasma directly)

- BBB is disrupted in brain traumatic, inflammatory or toxic inj., as well as in malignant brain tumors

→ influence choice and dose of antibiotics.

Blood-Brain Barrier (BBB)




② Blood-Cerebrospinal Fluid Barrier

Structure

- Endothelial cells with closed fenestrations
- Basement membrane
- Scattered pale cells
- Basement membrane
- Choroidal epithelial cells

➤ Tight junctions

not in vessels
but in the cuboidal cell



Cerebrospinal Fluid-Brain Interface


- A comparable physiologic barrier between CSF and the CNS extracellular compartment does not exist!

Clinical Notes


- تورم في قاع العين
• **Papilledema** *optic disc swelling in the eye base . [↑ pressure in the skull] ..retinal vein فينغ الدم مع الـ retinal vein*
- High ICP → high CSF pressure → Compression of retinal vein BL → optic disc swelling BL “Papilledema” → ON atrophy & blindness if persistent

- **Hydrocephalus** *and make symptoms*
- Abnormal increase in CSF formation (rare, like in CPP)
 - Blockage of CSF circulation (tumor, AS), noncommunicating HCP
 - ↓ CSF absorption (post meningitis, sinus thrombosis), communicating HCP

Hydrocephalus



normal Brain



acute hydrocephalus:
symptomatic.

CSF analysis in meningitis

← مسبب نفع ١ infection يغير التآيب لدر CSF ..

Tuberculosis	Fungal	Viral	Bacterial <i>most common</i>	Normal	درجه اعقاب
Variable	Elevated	Normal or slightly elevated	Elevated	5-20	Opening pressure cm H ₂ O
10 - 1000	10 - 1000	25 - 2000	> 1000	< 5	Leukocytes
Lymphocyte predominance	Lymphocyte predominance	Early: Neutrophils Late: Lymphocytes <i>تآيب</i>	Neutrophil predominance <i>not lymphocyte</i>	0 <i>and if it present mainly lymphocyte</i>	Cell differential
Very High	High	Normal or slightly elevated	High	15 - 45	Protein
Low	Low	Normal	Low <i>proteins or بشخصية او</i>	45 - 85	Glucose
+ (AFB)	+ (Fungal)	-	+	-	Cultures

lumbar puncture not for patient with tumor [because there is ↑ P]

4th ventricle tumors *common in children*

most common

- Medulloblastoma: arises from the vermis (roof) of the 4th V *من فوق النخاع*
Brain stem / نخاع *Cerebellum / مخ*

- Ependymoma: arises from the floor of the 4th V (can invade hypoglossal and vagal nuclei, affecting movements of the tongue and/or swallowing, respiration, heart rate, and blood pressure)
من تحت النخاع
Brain stem [floor of 4th ventricle]

4th ventricle tumor

coroid plexus papilloma

