

Physio Renal Dr. Hammam

1. In an asthmatic patient treated by B receptor agonist, potassium level was found to be 3.1 (normal 3.5-4.5). Which of the following is the mechanism by which this hypokalemia has developed?

- a) Increased renal excretion of potassium
- b) Increased renal secretion of potassium
- c) Transcellular shift of potassium
- d) Reduced gastrointestinal absorption of potassium
- e) Increased gastrointestinal absorption of potassium

2. In states of fluid deprivation, countercurrent mechanism helps developing high corticopapillary osmotic gradient in order to increase water reabsorption and induce

- a) an increase of plasma osmolarity
- b) a decrease of urine osmolarity
- c) an increase of aldosterone mediated potassium reabsorption
- d) an increase of urine osmolarity
- e) production of dilute urine

3. Two thirds (2/3) of total body water is confined to

- a) intracellular compartment
- b) Extracellular fluid compartment
- c) Intravascular compartment
- d) Interstitial compartment
- e) Plasma

4. In a patient with elevated (above normal) serum parathyroid hormone level, elevated serum calcium level develops secondary to

- a) Increased renal excretion of calcium
- b) Increased osteoclast function (bone resorption)
- c) Increased calcium reabsorption from the distal tubule and thick ascending loop of Henle
- d) Calcitonin-mediated increase of renal calcium reabsorption
- e) B&C

5. The main site of tubular magnesium reabsorption is

- a) Proximal tubule
- b) Glomerulus
- c) Collecting ducts
- d) Thick ascending loop of Henle
- e) Distal tubule

Physio Renal Dr. Hammam

6. Regarding phosphate renal handling, all the following is true, EXCEPT

- a) The main site of phosphate reabsorption in the nephron is the distal tubule
- b) 15% of the filtered phosphate is excreted
- c) PTH (parathyroid hormone) inhibits phosphate reabsorption
- d) 85% of the filtered phosphate is reabsorbed
- e) The main site of phosphate reabsorption in the nephron is proximal tubule

7. Regarding renal sodium handling, all the following is true EXCEPT

- a) Distal tubule is the main site of Na reabsorption in the nephron
- b) Na is reabsorbed along the entire nephron
- c) Na is reabsorbed in the proximal tubule by co-transport and counter-transport mechanisms
- d) Thick ascending limb of loop of Henle is sodium-permeable segment of the nephron
- e) Descending limb of the loop of Henle is sodium-impermeable segment of the nephron

8. Regarding the nerve supply of the urinary bladder and urethral sphincters, which of the following is TRUE?

- a) Sympathetic innervation represents the nerve of emptying
- b) Parasympathetic innervation represents the nerve of filling
- c) External urethral sphincter is innervated by autonomic nervous system
- d) Stimulation of parasympathetic nerve supply from S2, S3&S4 induces contraction of Detrusor muscle and relaxation of internal urethral sphincter
- e) Loss of function of higher inhibitory centers of micturition results in spastic neurogenic bladder

9. Regarding micturition reflex, all of the following is true EXCEPT

- a) It is self-regenerative
- b) It is elicited by stimulation of the stretch receptors in the urinary bladder wall
- c) It is facilitated by voluntary increase in the abdominal pressure
- d) Spinal centers of the reflex are L2, L3&L4
- e) It can be controlled voluntarily

- ✚ Volume contraction **increase** aldosterone & protein
- ✚ Volume expansion **decrease** aldosterone & protein
- ✚ Hyper-osmotic **increase** ADH & Na
- ✚ Iso-osmotic **no change** in ADH & Na
- ✚ Hypo-osmotic **decrease** Na
 - ❖ Contraction **no Change** in ADH
 - ❖ Expansion **decrease** ADH