

Question 4

Not yet
answered

Marked out of
1.0

🚩 Flag
question

The conscious production of signs and symptoms of a mental or physical illness, Unconscious motivation (without knowing why), No obvious external gains, most patients become angered when confronted and will leave hospital
Against medical advice

- ☐ a. malingering
- ☒ b. factitious disorder
- ☐ c. somatic symptom disorder
- ☐ d. conversion disorder
- ☐ e. none of the above

Question 5

Not yet
answered

Marked out of
1.0

🚩 Flag
question

Which of the following is a feature of the general adaptation syndrome described by Hans Selye

- ☐ a. resistance
- ☐ b. none of the answers
- ☐ c. exhaustion
- ☐ d. initial alarm
- ☒ e. all of the answers

Question 6

Not yet
answered

Marked out of
1.0

Flag
question

Which of the following is considered as a risk factor for suicide

- ☐ a. female gender
- ☐ b. marriage
- ☐ c. employment
- ☒ d. none of the answers
- ☐ e. all of the answers

Next page

All of the following mental disorders are common in males except one

- ☐ a. conduct disorder
- ☐ b. autism
- ☐ c. ADHD
- ☐ d. antisocial personality disorder
- ☒ e. Anorexia nervosa

Which of the following is the most common cause of intellectual disabilities

- ☐ a. fragile X syndrome
- ☐ b. autism
- ☒ c. fetal alcoholic syndrome
- ☐ d. down syndrome
- ☐ e. none of the answers

Male erectile disorder (impotence) has 10–20% lifetime prevalence; Half of men treated for sexual disorder complain of impotence. Which of the following sexual phases is affected by male erectile disorder (impotence)

- ☐ a. desire phase
- ☐ b. all of the answers
- ☒ c. arousal phase
- ☐ d. orgasmic phase

Question 10

Not yet
answered

Marked out of
1.0

Flag
question

When the patient is aware of the problem but ambivalent of starting treatment, what is the stage of behavioral change he is on

- ☐ a. relapse
- ☐ b. precontemplation
- ☐ c. preparation
- ☐ d. maintenance
- ☒ e. **contemplation**

Question 11

Not yet
answered

Marked out of
1.0

Flag
question

A 34-year-old secretary climbs 12 flights of stairs every day to reach her office because she is terrified by the thought of being trapped in the elevator. She has never had any traumatic event occur in an elevator; she has been terrified of them since childhood. Which of the following is the most likely diagnosis?

- ☐ a. Somatization disorder
- ☒ b. **specific phobia**
- ☐ c. malingering
- ☐ d. conversion disorder
- ☐ e. generalized anxiety disorder

Question 12

Not yet
answered

Marked out of
1.0

Flag
question

Which of the following is a feature of post-traumatic stress disorder

- ☐ a. re-experiencing the event
- ☐ b. mood instability
- ☐ c. increase startle reflex
- ☒ d. all of the answers
- ☐ e. phobic avoidance

Next page

Question 13

Not yet
answered

Marked out of
1.0

🚩 Flag
question

All of the followings are features of obsessions except one

- ☒ a. they are ego syntonic
- ☐ b. they are distressing
- ☐ c. they increases the anxiety level
- ☐ d. They are senseless
- ☐ e. they are time consuming

Question 14

Not yet
answered

Marked out of
1.0

🚩 Flag
question

A 27 years old female is referred to you by her family members, they stated that she has been married in the past ten years without any problems and she didn't have children, two months ago she decided to do (IVF, تلقيح مخبري), yesterday she received the result that the (IVF) failed, she didn't give any response, today morning she can't get out of her bed complaining of paralysis of both lower limbs. Which of the following is the most likely diagnosis

- ☐ a. factitious disorder
- ☐ b. malingering
- ☐ c. somatic symptom disorder
- ☐ d. none of the answers
- ☒ e. conversion disorder

Question 15

Not yet
answered

Marked out of
1.0

Flag
question

All of the followings are features of type (B) personality except one

- ☐ a. they don't mind losing
- ☐ b. they are not competitive
- ☐ c. they live in lower levels of stress
- ☒ d. they are impatient
- ☐ e. none of the answers

Next page

Question 16

Not yet
answered

Marked out of
1.0

🚩 Flag
question

All of the followings are causes of hypoactive sexual desire disorder in males except one

- ☐ a. marital discord
- ☒ b. none of the answers
- ☐ c. low testosterone level
- ☐ d. CNS depressant medications
- ☐ e. depression

Question 17

Not yet
answered

Marked out of
1.0

🚩 Flag
question

According to Holmes and Rahe scale used to quantify stressful life events. A man who takes 350 points on this scale that means that

- ☐ a. He had 50% chance of a major stress-induced health problem in next 2 years
- ☐ b. He had 100% chance of a major stress-induced health problem in next 2 years
- ☐ c. none of the above
- ☐ d. relatively low amount of life change and low susceptibility to stress-induced health problems
- ☒ e. He had 80% chance of a major stress-induced health problem in next 2 years

Question 18

Not yet
answered

Marked out of
1.0

Flag
question

A projective personality test in which the patient is asked to tell a story about what is going on in the picture

- ☐ a. projective drawing test
- ☒ b. Thematic apperception test
- ☐ c. sentence completion
- ☐ d. object assembly
- ☐ e. Rorschach inkblot test

Next page

Question 19

Not yet
answered

Marked out of
1.0

Flag
question

unable to conform to rules of society (only personality disorder that requires age 18 for diagnosis), Criminal acts: delinquency, theft, Unable to hold a job, Unable to maintain enduring attachments, Reckless, Aggressive, Show lack of remorse and more common in males than females

- ☐ a. narcissistic personality disorder
- ☒ b. antisocial personality disorder
- ☐ c. border line personality disorder
- ☐ d. histrionic personality disorder
- ☐ e. schizoid personality disorder

Question 20

Not yet
answered

Marked out of
1.0

Flag
question

Multiple motor and vocal tics that occur many times per day or intermittently for more than one year, Men > women with mean onset is age 7

- ☐ a. psychotic disorder
- ☐ b. ADHD
- ☐ c. conduct disorder
- ☐ d. obsessive compulsive disorder
- ☒ e. Tourette disorder

Question 21

Not yet
answered

Marked out of
1.0

Flag
question

Schizophrenia symptoms such as hallucinations and delusions PLUS mood disorder symptoms such as depression and mania which must have at least 2 weeks of psychotic symptoms in the absence of mood symptoms, this is called

- ☐ a. Schizotypal disorder
- ☐ b. schizoid disorder
- ☒ c. schizoaffective disorder
- ☐ d. schizophrenia
- ☐ e. schizophreniform disorder

Next page

Question 22

Not yet
answered

Marked out of
1.0

Flag
question

It has been observed that schizophrenia is more common between monozygotic twins than dizygotic twin, this observation goes with which of the following theories of schizophrenia development

- ☐ a. viral theory
- ☐ b. social theory
- ☒ c. genetic theory
- ☐ d. biological theory
- ☐ e. none of the answers

Question 23

Not yet
answered

Marked out of
1.0

Flag
question

All of the followings are features of schizophrenia except one

- ☒ a. lasted less than one month
- ☐ b. disorganized speech
- ☐ c. hallucinations
- ☐ d. negative symptoms
- ☐ e. delusions

Question 24

Not yet
answered

Marked out of
1.0

Flag
question

A 28-year-old man referred to a psychiatrist for evaluation, in the past six months he used to experience multiple attacks of (chest pain, dyspnea, palpitation, tremors, fainting and he feels that he will loss control and die) these attacks lasts about ten minutes each, multiple investigations done to be normal. What is your diagnosis?

- ☐ a. factitious disorder
- ☐ b. Somatization disorder
- ☒ c. panic attacks
- ☐ d. conversion disorder
- ☐ e. malingering

Next page

Question 25

Not yet
answered

Marked out of
1.0

Flag
question

Grandiose sense of self-importance, Preoccupied with fantasies of unlimited wealth, power, love, Demands constant attention and admiration and meets criticism with indifference or rage all these are features of

- ☐ a. histrionic personality disorder
- ☒ b. narcissistic personality disorder
- ☐ c. antisocial personality disorder
- ☐ d. border line personality disorder
- ☐ e. schizoid personality disorder

Question 26

Not yet
answered

Marked out of
1.0

Flag
question

A 32-year-old woman is brought to the emergency room when she complains of chest pain. She is noted to be hyper vigilant and anxious, with a pulse of 120 beats per minute and BP of 160/97. She has widely dilated pupils. Her toxicology screen is positive. Which of the following drugs is she most likely to have used?

- ☐ a. Benzodiazepines
- ☐ b. Cannabis
- ☒ c. Cocaine
- ☐ d. morphine
- ☐ e. Heroin

Question 27

Not yet
answered

Marked out of
1.0

Flag
question

Involves sudden unexpected travel, an inability to recall one's past, or confusion of identity

- ☐ a. depersonalization
- ☐ b. dissociative identity disorder
- ☐ c. derealization
- ☐ d. dissociative personality disorder
- ☒ e. dissociative fugue disorder

Next page

Question 28

Not yet
answered

Marked out of
1.0

🚩 Flag
question

All of the followings are features of odd or eccentric personality disorders except one

- ☐ a. they have peculiar ways of thinking
- ☒ b. they are dramatic and emotionally laden
- ☐ c. They have odd behaviors
- ☐ d. they are socially withdrawn
- ☐ e. they are suspicious of others

Question 29

Not yet
answered

Marked out of
1.0

🚩 Flag
question

Age of onset of schizophrenia between males is

- ☐ a. 45 -- 55 years
- ☐ b. 25 -- 35 years
- ☐ c. 55 -- 65 years
- ☐ d. 35 – 45 years
- ☒ e. 15 – 25 years

Question 30

Not yet
answered

Marked out of
1.0

Flag
question

In Paraphilic Disorders a sexual pleasure derived from others' pain to achieve orgasm is called

- ☐ a. Exhibitionism
- ☐ b. Pedophilia
- ☐ c. Masochism
- ☐ d. Frotteurism
- ☒ e. Sadism

Finish attempt ...