

Pharma

Dr. AM Fouda, Professor of Clinical Pharmacology

Summaries of Ali Al-Murtada

ملخصات

الدكتور عبد المتعال فودة

أعداد: الطالب علي المرتضى



Anti-depressant drugs

- 5% of People have depression
- Women > Men (2:1), Major disability

Causes :-

- 1- 4 genes
- 2- Biogenic ~~and~~ amine and receptor theory

NA ↓, Serotonin ↓, Dopamine ↓
زيادة في α -5HT_{2A}
 β -5HT_{2C}

- 3- ~~Neurotropic~~ Neurotropic and Cytokines therapy
 - A - Brain derived neurotrophic factor (BDNF) ↓
 - B - Hypothalamus - Pituitary axis dysregulation
 - C - Proinflammatory Cytokines = IL-1, IL-6, TNF α

Types :- 1- Unipolar most common

2- Bipolar نادر

Symptoms :-

A- Emotional :- Serotonin ↓

- 1- Anhedonia :- lose interest and enjoyment
- 2- Apathy and loss of self-esteem
- 3- loss of motivation

B- Biological :- NA ↓

- 1- Sleep disturbance and loss appetite
- 2- loss libido
- 3- **Chronic Pain**

Drugs :-

- 1- Monoamine oxidase inhibitors (MAOIs) لا يحتمل
- 2- Tricyclic and Tetracyclic anti-depressant (TCA)
- 3- Selective serotonin reuptake inhibitor (SSRIs)
- 4- Atypical (heterocyclic) antidepressant → شائع

MAOIs → **dopamine only**

- MAOA in Brain and GIT, MAOB Brain only (liver)
- Drugs : Phenelzine, Tranylcypromine, **Sedgiline**

Mechanism :- ~~MAO~~ → NA ↑

also Parkinson

Uses:

- 1- depression that unrelieved by other drugs
- 2- Parkinsonism: Selegiline (MAOB only) $\rightarrow$ Dopamine $\uparrow$
Selective

Adverse effect أول أربعة آثار

- 1- Orthostatic hypotension (postural) $\rightarrow$ أي علاج يغير بالدماغ يسي hypo (X₂)
- 2- Sexual dysfunction
- 3- Serotonin Syndrome, if mix with TCR and SSRIs
- 4- CNS stimulation $\rightarrow$ irritability, insomnia, tremor, hyperthermia
Convulsions
- 5- Chesse reaction (hypertensive crisis) occur when eating
Tyramine-rich food eg: yogurt, old chesse
أين عم NA
Broke by MAO in liver

TCA

Drugs: Imipramine, Clomiramine, Ami tripty line, Nortriptylines

Mechanism: inhibits of NA and Serotonin by block transporter that responsible for it (inhibits amine pump)

Uses: 1- depression
2- Chronic Pain Condition eg. Fibromyalgia
3- Nocturnal enuresis in children (bed-wetting) use imipramine

Adverse effect أول أعراضه النفسية

- 5- CNS Sedation (atropine-like action)
- 6- Atropine-like action
- 7- Cardiac arrhythmia
- 8- Hepatotoxicity: cholestatic jaundice

أول أعراضه

TCA Toxicity

Manifestations :

- 1- Atropine-like action
- 2- Metabolic acidosis
- 3- Cardiac arrhythmia & Tachycardia, Wide QRS, long QT

Treatment :

- 1- I.V NaCO_3 : 1st line (هذا ممكن كافي)
- 2- I.V lidocaine

* dialysis not effective

SSRIs صاروخ

Drugs :- Fluoxetine, Paroxetine, Sertraline, Citalopram, escitalopram

Uses :

- 1- Major depression
- 2- OCD الوسواس القهري
- 3- General anxiety disorder (GAD)

Adverse effect :- آثار الأضرار

- 4- CNS : sedation or excitation gone with time
 - 5- GIT irritation (Common) :- Prescribed with PPIs
 - 6- Cardiac arrhythmia :- QT interval prolongation
 - * Sertraline (safe) :- Treat depression after MI
 - * SSRIs :- stop gradually over 4 weeks
-

Atypical الأتypical

A- NA and 5HT reuptake inhibitors also block 5HT_{2A} and 5HT_{2C}

- Venlafaxine
- Duloxetine : used more for autonomic neuropathy

B- Receptor blockers :-

- Trazodone : Block 5HT_{2A} and 5HT_{2C} and more sedative
- Mirtazapine : Block α_2 and 5HT_{2C}, less GIT upset and sexual dysfunction than SSRIs